

# ABALOPARATIDE

---

## Products Affected

- TYMLOS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 MONTHS
<b>Other Criteria</b>	OSTEOPOROSIS: HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ABATACEPT IV

---

## Products Affected

- ORENCIA (WITH MALTOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
<b>Coverage Duration</b>	RA, PJIA, PSA: INITIAL: 6 MOS, RENEWAL: 12 MOS. ACUTE GRAFT VERSUS HOST DISEASE (AGVHD): 1 MO.
<b>Other Criteria</b>	INITIAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. RENEWAL: RA, PJIA, PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ABATACEPT SQ

---

## Products Affected

- ORENCIA
- ORENCIA CLICKJECT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# ABEMACICLIB

---

## Products Affected

- VERZENIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ABIRATERONE

---

## Products Affected

- *abiraterone*
- *abirtega*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC), METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ABIRATERONE SUBMICRONIZED

---

## Products Affected

- *abiraterone, submicronized*
- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ACALABRUTINIB

---

## Products Affected

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ACORAMIDIS

## Products Affected

- ATTRUBY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CARDIOMYOPATHY OF WILD TYPE OR VARIANT TRANSTHYRETIN-MEDIATED AMYLOIDOSIS (ATTR-CM): INITIAL: 1) NEW YORK HEART ASSOCIATION (NYHA) CLASS I, II, OR III HEART FAILURE, AND 2) DIAGNOSIS CONFIRMED BY (A) BONE SCAN (SCINTIGRAPHY) STRONGLY POSITIVE FOR MYOCARDIAL UPTAKE OF TC-99M-PYP, OR (B) BIOPSY OF TISSUE OF AFFECTED ORGAN(S) (CARDIAC AND POSSIBLY NON-CARDIAC SITES) TO CONFIRM AMYLOID PRESENCE AND CHEMICAL TYPING TO CONFIRM PRESENCE OF TRANSTHYRETIN (TTR) PROTEIN.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ATTR-CM: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ATTR SPECIALIST, OR MEDICAL GENETICIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ATTR-CM: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER ATTR-CM TTR STABILIZERS (E.G., TAFAMIDIS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ADAGRASIB

---

## Products Affected

- KRAZATI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ADALIMUMAB

## Products Affected

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UVEITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)
- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST.
<b>Coverage Duration</b>	INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. UVEITIS: DOES NOT HAVE ISOLATED ANTERIOR UVEITIS. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), PSA, AS, PSO, HS, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ADALIMUMAB-AATY

---

## Products Affected

- YUFLYMA(CF)
- YUFLYMA(CF) AI CROHN'S-UC-HS
- YUFLYMA(CF) AUTOINJECTOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST.
<b>Coverage Duration</b>	INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. UVEITIS: DOES NOT HAVE ISOLATED ANTERIOR UVEITIS. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), PSA, AS, PSO, HS, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ADALIMUMAB-ADBIM

## Products Affected

- CYLTEZO(CF)
- CYLTEZO(CF) PEN
- CYLTEZO(CF) PEN CROHN'S-UC-HS
- CYLTEZO(CF) PEN PSORIASIS-UV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST.
<b>Coverage Duration</b>	INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. UVEITIS: DOES NOT HAVE ISOLATED ANTERIOR UVEITIS. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), PSA, AS, PSO, HS, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# AFATINIB

---

## Products Affected

- GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ALECTINIB

---

## Products Affected

- ALECENSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ALPELISIB-PIQRAY

---

## Products Affected

- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AMIKACIN LIPOSOMAL INH

## Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MYCOBACTERIUM AVIUM COMPLEX (MAC) LUNG DISEASE: RENEWAL: 1) NO POSITIVE MAC SPUTUM CULTURE AFTER CONSECUTIVE NEGATIVE CULTURES, AND 2) IMPROVEMENT IN SYMPTOMS. ADDITIONALLY, FOR FIRST RENEWAL, APPROVAL REQUIRES AT LEAST ONE NEGATIVE SPUTUM CULTURE FOR MAC BY SIX MONTHS OF ARIKAYCE TREATMENT. FOR SECOND AND SUBSEQUENT RENEWALS, APPROVAL REQUIRES AT LEAST THREE NEGATIVE SPUTUM CULTURES FOR MAC BY 12 MONTHS OF ARIKAYCE TREATMENT.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	MAC LUNG DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 6 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AMIVANTAMAB-HYALURONIDASE-LPUJ

---

## Products Affected

- RYBREVANT FASPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AMIVANTAMAB-VMJW

---

## Products Affected

- RYBREVANT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ANAKINRA

## Products Affected

- KINERET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
<b>Required Medical Information</b>	INITIAL: CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR S100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES. DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE IL1RN GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR), AND 2) ONE OF THE FOLLOWING: PUSTULAR PSORIASIS-LIKE RASHES, OSTEOMYELITIS, ABSENCE OF BACTERIAL OSTEOMYELITIS, ONYCHOMADESIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. CAPS, DIRA: LIFETIME.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: RA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# APALUTAMIDE

---

## Products Affected

- ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC), METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# APOMORPHINE - ONAPGO

---

## Products Affected

- ONAPGO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PD: RENEWAL: IMPROVEMENT IN MOTOR SYMPTOMS WHILE ON THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# APOMORPHINE - SL

---

## Products Affected

- KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# APREMILAST

---

## Products Affected

- OTEZLA
- OTEZLA STARTER
- OTEZLA XR
- OTEZLA XR INITIATION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: MILD PLAQUE PSORIASIS (PSO): 1) PSORIASIS COVERING LESS THAN 3 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC THERAPY (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) OR ONE CONVENTIONAL TOPICAL THERAPY (E.G., TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. BEHCETS DISEASE: 1) HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: ALL INDICATIONS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ARIMOCLOMOL

---

## Products Affected

- MIPLYFFA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NIEMANN-PICK DISEASE TYPE C (NPC): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH NEUROLOGIST OR GENETICIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	NPC: RENEWAL: IMPROVEMENT OR SLOWING OF DISEASE PROGRESSION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ASCIMINIB

---

## Products Affected

- SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PREVIOUSLY TREATED OR T315I MUTATION PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ASFOTASE ALFA

---

## Products Affected

- STRENSIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6 MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II) CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V) NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM CALCIUM, (VI) HISTORY OR PRESENCE OF NON-TRAUMATIC POSTNATAL FRACTURE AND DELAYED FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC DEFORMITIES, (II) PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5 YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OR PRESENCE OF NON-TRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3)</p>
	<p>NOT HAVE A TREATABLE FORM OF RICKETS. RENEWAL: ALL INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ATOGEPANT

---

## Products Affected

- QULIPTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AVACOPAN

---

## Products Affected

- TAVNEOS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ANTI-NEUTROPHIL CYTOPLASMIC AUTOANTIBODY (ANCA)-ASSOCIATED VASCULITIS: INITIAL: ANCA SEROPOSITIVE (ANTI-PR3 OR ANTI-MPO).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ANCA-ASSOCIATED VASCULITIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 6 MONTHS.
<b>Other Criteria</b>	ANCA-ASSOCIATED VASCULITIS: RENEWAL: CONTINUES TO BENEFIT FROM THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AVAPRITINIB

---

## Products Affected

- AYVAKIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AVUTOMETINIB-DEFACTINIB

---

## Products Affected

- AVMAPKI
- AVMAPKI-FAKZYNJA
- FAKZYNJA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AXATILIMAB-CSFR

---

## Products Affected

- NIKTIMVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CHRONIC GRAFT VS HOST DISEASE (CGVHD): 1) FAILURE OF AT LEAST TWO LINES OF SYSTEMIC THERAPY, ONE OF WHICH MUST BE A TRIAL OF OR CONTRAINDICATION TO JAKAFI, AND 2) NO CONCURRENT USE WITH JAKAFI, REZUROCK, OR IMBRUVICA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# AXITINIB

---

## Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AZACITIDINE

---

## Products Affected

- ONUREG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AZTREONAM INHALED

---

## Products Affected

- CAYSTON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	7 YEARS OF AGE OR OLDER
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BEDAQUILINE

---

## Products Affected

- SIRTURO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 WEEKS
<b>Other Criteria</b>	N/A
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BELIMUMAB

## Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BELUMOSUDIL

---

## Products Affected

- REZUROCK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CHRONIC GRAFT VS HOST DISEASE (CGVHD): 1) FAILURE OF AT LEAST TWO LINES OF SYSTEMIC THERAPY, ONE OF WHICH MUST BE A TRIAL OF OR CONTRAINDICATION TO JAKAFI, AND 2) NO CONCURRENT USE WITH JAKAFI, NIKTIMVO, OR IMBRUVICA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# BELZUTIFAN

---

## Products Affected

- WELIREG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BENDAMUSTINE

---

## Products Affected

- *bendamustine intravenous recon soln*
- BENDAMUSTINE INTRAVENOUS SOLUTION
- BENDEKA
- VIVIMUSTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# BENRALIZUMAB

---

## Products Affected

- FASENRA
- FASENRA PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING AT LEAST 3 DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: ASTHMA: 1) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 2) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS (EGPA): REDUCTION IN EGPA SYMPTOMS COMPARED TO BASELINE OR ABILITY TO REDUCE/ELIMINATE CORTICOSTEROID USE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BETAINE

---

## Products Affected

- *betaine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# BEVACIZUMAB-BVZR

---

## Products Affected

- ZIRABEV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BEXAROTENE

---

## Products Affected

- *bexarotene*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BINIMETINIB

---

## Products Affected

- MEKTOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BORTEZOMIB

---

## Products Affected

- *bortezomib injection*
- BORUZU

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BOSENTAN

## Products Affected

- *bosentan oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NO CONCURRENT USE WITH CYCLOSPORINE A OR GLYBURIDE. RENEWAL: NO CONCURRENT USE WITH CYCLOSPORINE A OR GLYBURIDE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# BOSUTINIB

## Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# BRIGATINIB

---

## Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# C1 ESTERASE INHIBITOR-HAEGARDA

## Products Affected

- HAEGARDA SUBCUTANEOUS RECON SOLN 2,000 UNIT, 3,000 UNIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY ANGIOEDEMA (HAE): INITIAL: 1) TYPE III HAE, OR 2) TYPE I OR II HAE CONFIRMED BY ONE OF THE FOLLOWING COMPLEMENT TESTS: C1-INH PROTEIN LEVELS, C4 PROTEIN LEVELS, C1-INH FUNCTIONAL LEVELS, C1Q.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, ALLERGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CABOZANTINIB CAPSULE

---

## Products Affected

- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CABOZANTINIB TABLET

---

## Products Affected

- CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CANNABIDIOL

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CAPIVASERTIB

---

## Products Affected

- TRUQAP

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CAPMATINIB

---

## Products Affected

- TABRECTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CARGLUMIC ACID

## Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ACUTE OR CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.
<b>Other Criteria</b>	RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CERITINIB

---

## Products Affected

- ZYKADIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No



PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, HUMIRA/CYLTEZO/YUFLYMA, SELARSDI/YESINTEK, XELJANZ, RINVOQ, SKYRIZI, TREMFYA, ORENCIA, OTEZLA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: SELARSDI/YESINTEK, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, HUMIRA/CYLTEZO/YUFLYMA, SELARSDI/YESINTEK, SKYRIZI, TREMFYA, OTEZLA. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ. INITIAL FOR RA, PSA, PSO, AS, CD, PJIA: TRIAL OF OR CONTRAINDICATION TO THE STEP AGENTS IS NOT REQUIRED IF THE PATIENT IS PREGNANT, BREASTFEEDING, OR TRYING TO BECOME PREGNANT. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL FOR RA, PSA, AS, PSO, NR-AXSPA, PJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CETUXIMAB

---

## Products Affected

- ERBITUX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CLADRIBINE

## Products Affected

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	48 WEEKS.
<b>Other Criteria</b>	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CLOBAZAM-SYMPAZAN

---

## Products Affected

- SYMPAZAN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	LGS: INITIAL: CONTRAINDICATION TO OR UNABLE TO SWALLOW CLOBAZAM TABLETS OR SUSPENSION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# COBIMETINIB

---

## Products Affected

- COTELLIC

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CORTICOTROPIN

## Products Affected

- CORTROPHIN GEL INJECTION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.
<b>Coverage Duration</b>	INFANTILE SPASMS AND MS: 28 DAYS. ALL OTHER FDA APPROVED INDICATIONS: INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA-PD PLAN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	Yes

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# CRIZOTINIB CAPSULE

---

## Products Affected

- XALKORI ORAL CAPSULE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CRIZOTINIB PELLETS

---

## Products Affected

- XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-SMALL CELL LUNG CANCER (NSCLC), ANAPLASTIC LARGE CELL LYMPHOMA (ALCL), INFLAMMATORY MYOFIBROBLASTIC TUMOR (IMT): UNABLE TO SWALLOW CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DABRAFENIB CAPSULES

---

## Products Affected

- TAFINLAR ORAL CAPSULE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DABRAFENIB SUSPENSION

---

## Products Affected

- TAFINLAR ORAL TABLET FOR SUSPENSION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNABLE TO SWALLOW TAFINLAR CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DACOMITINIB

---

## Products Affected

- VIZIMPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DALFAMPRIDINE

---

## Products Affected

- *dalfampridine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY (E.G., MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS, UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA). RENEWAL: IMPROVEMENT IN WALKING ABILITY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DAROLUTAMIDE

## Products Affected

- NUBEQA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC), METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DASATINIB

---

## Products Affected

- *dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND DASATINIB IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DATOPOTAMAB DERUXTECAN-DLNK

---

## Products Affected

- DATROWAY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DECITABINE/CEDAZURIDINE

---

## Products Affected

- INQOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DEFERASIROX

## Products Affected

- *deferasirox oral granules in packet*
- *deferasirox oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L. CHRONIC IRON OVERLOAD IN NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS), AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G OF LIVER DRY WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L. NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G OF LIVER DRY WEIGHT OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS OR NTDT: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS OR NTDT: DEFERASIROX SPRINKLE PACKETS: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX ORAL TABLET OR TABLET FOR ORAL SUSPENSION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DENOSUMAB-BMWO - OSENVELT

---

## Products Affected

- OSENVELT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DENOSUMAB-XGEVA

---

## Products Affected

- XGEVA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DEUTETRABENAZINE

## Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 18 MG, 6 MG
- AUSTEDO XR TITRATION KT(WK1-4)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TARDIVE DYSKINESIA: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DICLOFENAC TOPICAL SOLUTION

---

## Products Affected

- *diclofenac sodium topical solution in metered-dose pump*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# DICLOFENAC-FLECTOR

---

## Products Affected

- *diclofenac epolamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# DIMETHYL FUMARATE

---

## Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# DIROXIMEL FUMARATE

---

## Products Affected

- VUMERITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DORDAVIPRONE

---

## Products Affected

- MODEYSO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DOSTARLIMAB-GXLY

---

## Products Affected

- JEMPERLI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DRONABINOL CAPSULE

---

## Products Affected

- *dronabinol*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 MONTHS
<b>Other Criteria</b>	NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DROXIDOPA

## Products Affected

- *droxidopa*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
<b>Coverage Duration</b>	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
<b>Other Criteria</b>	NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DUPILUMAB

## Products Affected

- DUPIXENT PEN
- DUPIXENT SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL OF 150 TO 1500 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY. ATOPIC DERMATITIS (AD): AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: AD, PN, CSU: PRESCRIBED OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED OR IN CONSULTATION WITH ALLERGIST OR PULMONOLOGIST. CRSWNP: PRESCRIBED OR IN CONSULTATION WITH OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED OR IN CONSULTATION WITH GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST. COPD: PRESCRIBED OR IN CONSULTATION WITH PULMONOLOGIST. RENEWAL: CSU: PRESCRIBED OR IN CONSULTATION WITH ALLERGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	BP: 12 MO. AD/CRSWNP/EOE/PN/CSU: INITIAL/RENEWAL: 6 MO/12 MO. ASTHMA/COPD: INITIAL/RENEWAL: 12 MO.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: AD: 1) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR). ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING AT LEAST 3 DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA. CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP): 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, 2) EVIDENCE OF NASAL POLYPS BY DIRECT EXAMINATION, ENDOSCOPY, OR SINUS CT SCAN, AND 3) INADEQUATELY CONTROLLED DISEASE. PRURIGO NODULARIS (PN): CHRONIC PRURITUS (ITCH MORE THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, AND HISTORY OR SIGN OF A PROLONGED SCRATCHING BEHAVIOR. EOSINOPHILIC COPD: USED IN COMBINATION WITH A LAMA/LABA/ICS. CHRONIC SPONTANEOUS URTICARIA (CSU): 1) TRIAL OF AND MAINTAINED ON, OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE AND 2) STILL EXPERIENCES HIVES OR ANGIOEDEMA MOST DAYS OF THE WEEK FOR AT LEAST 6 WEEKS. INITIAL/RENEWAL : ALL INDICATIONS EXCEPT BULLOUS PEMPHIGOID (BP): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4</p>

PA Criteria	Criteria Details
	<p>INHIBITOR) FOR THE SAME INDICATION. RENEWAL: AD, CRSWNP, EOE: IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 2) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITUS OR PRURIGINOUS LESIONS. EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A LAMA/LABA/ICS, AND 2) CLINICAL RESPONSE AS EVIDENCED BY (A) REDUCTION IN COPD EXACERBATIONS FROM BASELINE, (B) REDUCTION IN SEVERITY OR FREQUENCY OF COPD-RELATED SYMPTOMS, OR (C) INCREASE IN FEV1 OF AT LEAST 5 PERCENT FROM PRETREATMENT BASELINE. CSU: MAINTAINED ON OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# DUVELISIB

---

## Products Affected

- COPIKTRA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# EFLORNITHINE

---

## Products Affected

- IWILFIN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ELACESTRANT

---

## Products Affected

- ORSERDU ORAL TABLET 345 MG, 86 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ELAGOLIX

## Products Affected

- ORLISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.
<b>Prescriber Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND A PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# ELAPEGADEMASE-LVLR

## Products Affected

- REVC0VI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ADENOSINE DEAMINASE SEVERE COMBINED IMMUNE DEFICIENCY (ADA-SCID): INITIAL: ADA-SCID AS MANIFESTED BY: 1) CONFIRMATORY GENETIC TEST, OR 2) SUGGESTIVE LABORATORY FINDINGS (E.G., ELEVATED DEOXYADENOSINE NUCLEOTIDE [DAXP] LEVELS, LYMPHOPENIA) AND HALLMARK SIGNS/SYMPTOMS (E.G., RECURRENT INFECTIONS, FAILURE TO THRIVE, PERSISTENT DIARRHEA).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ADA-SCID: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH IMMUNOLOGIST, HEMATOLOGIST/ONCOLOGIST, OR PHYSICIAN SPECIALIZING IN INHERITED METABOLIC DISORDERS.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ADA-SCID: RENEWAL: 1) IMPROVEMENT OR MAINTENANCE OF IMMUNE FUNCTION FROM BASELINE, AND 2) HAS NOT RECEIVED SUCCESSFUL HCT OR GENE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ELEXACAFITOR-TEZACAFITOR-IVACAFITOR

## Products Affected

- TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL
- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CYSTIC FIBROSIS (CF): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
<b>Other Criteria</b>	CF: INITIAL: NO CONCURRENT USE WITH ANOTHER CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR (CFTR) MODULATOR. RENEWAL: 1) IMPROVEMENT IN CLINICAL STATUS, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ELRANATAMAB-BCMM

## Products Affected

- ELREXFIO 44 MG/1.1 ML VIAL INNER, SUV, P/F
- ELREXFIO SUBCUTANEOUS SOLUTION 40 MG/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	RELAPSED OR REFRACTORY MULTIPLE MYELOMA: RENEWAL: 1) HAS RECEIVED AT LEAST 24 WEEKS OF TREATMENT WITH ELREXFIO, AND 2) HAS RESPONDED TO TREATMENT (PARTIAL RESPONSE OR BETTER), AND HAS MAINTAINED THIS RESPONSE FOR AT LEAST 2 MONTHS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ELTROMBOPAG - ALVAIZ

## Products Affected

- ALVAIZ

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: 1) PLATELET COUNT IS LESS THAN $30 \times 10^9/L$ , OR 2) PLATELET COUNT IS LESS THAN $50 \times 10^9/L$ AND HAD A PRIOR BLEEDING EVENT.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
<b>Coverage Duration</b>	ITP: INITIAL: 6 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
<b>Other Criteria</b>	INITIAL: ITP: 1) TRIAL OF OR CONTRAINDICATION TO ONE CORTICOSTEROID OR IMMUNOGLOBULIN, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY, AND 2) NO CONCURRENT USE WITH OTHER THROMBOPOIETIN RECEPTOR AGONISTS (TPO-RAS). RENEWAL: ITP: 1) IMPROVEMENT IN PLATELET COUNT FROM BASELINE OR REDUCTION IN BLEEDING EVENTS, AND 2) NO CONCURRENT USE WITH OTHER TPO-RAS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ELTROMBOPAG - PROMACTA

## Products Affected

- *eltrombopag olamine oral powder in packet*  
12.5 mg, 25 mg
- *eltrombopag olamine oral tablet* 12.5 mg, 25 mg, 50 mg, 75 mg

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: 1) PLATELET COUNT OF LESS THAN $30 \times 10^9/L$ , OR 2) PLATELET COUNT OF LESS THAN $50 \times 10^9/L$ AND A PRIOR BLEEDING EVENT.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
<b>Coverage Duration</b>	ITP: INITIAL: 6 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
<b>Other Criteria</b>	INITIAL: ITP: 1) TRIAL OF OR CONTRAINDICATION TO ONE CORTICOSTEROID OR IMMUNOGLOBULIN, OR HAD AN INSUFFICIENT RESPONSE TO SPLENECTOMY, AND 2) NO CONCURRENT USE WITH OTHER THROMBOPOIETIN RECEPTOR AGONISTS (TPO-RAS). ALL INDICATIONS: ELTROMBOPAG ORAL SUSPENSION PACKETS: TRIAL OF A FORMULARY VERSION OF ELTROMBOPAG TABLET OR PATIENT IS UNABLE TO TOLERATE TABLET FORMULATION. RENEWAL: ITP: 1) IMPROVEMENT IN PLATELET COUNTS FROM BASELINE OR REDUCTION IN BLEEDING EVENTS, AND 2) NO CONCURRENT USE WITH OTHER TPO-RAS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# ENASIDENIB

---

## Products Affected

- IDHIFA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ENCORAFENIB

---

## Products Affected

- BRAFTOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ENSARTINIB

---

## Products Affected

- ENSACOVE ORAL CAPSULE 100 MG, 25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ENTRECTINIB CAPSULES

---

## Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ENTRECTINIB PELLETS

## Products Affected

- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), SOLID TUMORS: 1) TRIAL OF OR CONTRAINDICATION TO ROZLYTREK CAPSULES MADE INTO AN ORAL SUSPENSION, AND 2) DIFFICULTY OR UNABLE TO SWALLOW CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ENZALUTAMIDE

## Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (NMCSPC): HIGH RISK FOR METASTASIS (I.E. PSA DOUBLING TIME OF 9 MONTHS OR LESS). METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC), NON-METASTATIC CRPC (NMCRPC), METASTATIC CSPC (MCSPC), NMCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# EPCORITAMAB-BYSP

---

## Products Affected

- EPKINLY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# EPOETIN ALFA-EPBX

## Products Affected

- RETACRIT INJECTION SOLUTION  
10,000 UNIT/ML, 2,000 UNIT/ML, 20,000  
UNIT/2 ML, 20,000 UNIT/ML, 3,000  
UNIT/ML, 4,000 UNIT/ML, 40,000  
UNIT/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL IS LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL IS 13G/DL OR LESS. RENEWAL: 1) CKD IN ADULTS NOT ON DIALYSIS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND THE DOSE IS BEING OR HAS BEEN REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. 2) CKD IN PEDIATRIC PATIENTS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS APPROACHED OR EXCEEDS 12G/DL AND THE DOSE IS BEING OR HAS BEEN REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. 3) ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. 4) CANCER CHEMOTHERAPY: (A) HEMOGLOBIN LEVEL IS LESS THAN 10 G/DL, OR (B) HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.
<b>Other Criteria</b>	RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ERDAFITINIB

---

## Products Affected

- BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ERENUMAB-AOOE

---

## Products Affected

- AIMOVIG AUTOINJECTOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ERLOTINIB

---

## Products Affected

- *erlotinib oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ESKETAMINE

## Products Affected

- SPRAVATO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST OR OTHER REMS-CERTIFIED PROVIDER.
<b>Coverage Duration</b>	INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: TRD, MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ETANERCEPT

---

## Products Affected

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION.</p> <p>INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: ALL INDICATIONS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# EVEROLIMUS-AFINITOR

---

## Products Affected

- *everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *torpenz oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# EVEROLIMUS-AFINITOR DISPERZ

---

## Products Affected

- *everolimus (antineoplastic) oral tablet for suspension*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# FECAL MICROBIOTA CAPSULE

## Products Affected

- VOWST

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	30 DAYS
<b>Other Criteria</b>	CLOSTRIDIoidES DIFFICILE INFECTION (CDI): 1) HAS NOT PREVIOUSLY RECEIVED VOWST: COMPLETION OF ANTIBIOTIC TREATMENT FOR RECURRENT CDI (AT LEAST 3 CDI EPISODES), OR 2) PREVIOUSLY RECEIVED VOWST: (A) TREATMENT FAILURE (DEFINED AS THE PRESENCE OF CDI DIARRHEA WITHIN 8 WEEKS OF FIRST DOSE OF VOWST AND A POSITIVE STOOL TEST FOR C. DIFFICILE), AND (B) HAS NOT RECEIVED MORE THAN ONE TREATMENT COURSE OF VOWST WHICH WAS AT LEAST 12 DAYS AND NOT MORE THAN 8 WEEKS PRIOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FEDRATINIB

---

## Products Affected

- INREBIC

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# FENFLURAMINE

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS); PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# FENTANYL CITRATE

---

## Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# FEZOLINETANT

## Products Affected

- VEOZAH

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MENOPAUSAL VASOMOTOR SYMPTOMS (VMS): INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO HORMONAL THERAPY (E.G., ESTRADIOL TRANSDERMAL PATCH, ORAL CONJUGATED ESTROGENS), 2) LABORATORY TESTING TO ESTABLISH BASELINE HEPATIC FUNCTION AND CONTINUED MONITORING OF THESE VALUES IN ACCORDANCE WITH THE FDA CURRENT LABEL RECOMMENDATION, AND 3) NO CONCURRENT USE WITH ANOTHER HORMONAL (E.G., PREMPRO) OR NON-HORMONAL (E.G., BRISDELLE) AGENT FOR VMS. RENEWAL: 1) CONTINUED NEED FOR VMS TREATMENT (PERSISTENT HOT FLASHES), 2) REDUCTION IN VMS FREQUENCY OR SEVERITY DUE TO VEOZAH TREATMENT, AND 3) NO NEW SYMPTOMS OF LIVER INJURY AND/OR WORSENING LAB VALUES (E.G., ALT, AST, TOTAL BILIRUBIN).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# FILGRASTIM-AAFI

---

## Products Affected

- NIVESTYM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FINERENONE

## Products Affected

- KERENDIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: HEART FAILURE (HF): 1) NEW YORK HEART ASSOCIATION (NYHA) CLASS II-IV, AND 2) LEFT VENTRICULAR EJECTION FRACTION OF AT LEAST 40 PERCENT NOT DUE TO AN UNDERLYING CAUSE (E.G., INFILTRATIVE CARDIOMYOPATHY, HYPERTROPHIC CARDIOMYOPATHY, VALVULAR DISEASE, PERICARDIAL DISEASE, HIGH-OUTPUT HEART FAILURE).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: HF: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	CHRONIC KIDNEY DISEASE (CKD) ASSOCIATED WITH TYPE 2 DIABETES (T2D): INITIAL: HISTORY OF AND WILL CONTINUE ON, HAS A CONTRAINDICATION, OR INTOLERANCE TO AN ANGIOTENSIN CONVERTING ENZYME INHIBITOR (ACE-I) OR AN ANGIOTENSIN RECEPTOR BLOCKER (ARB). HF: INITIAL/RENEWAL: NO CONCURRENT USE WITH ANOTHER MINERALOCORTICOID (ALDOSTERONE) RECEPTOR ANTAGONIST.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# FINGOLIMOD

---

## Products Affected

- *fingolimod*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# FOSCARBIDOPA-FOSLEVODOPA

## Products Affected

- VYALEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PD: INITIAL: ONE OF THE FOLLOWING: 1) UNABLE TO SWALLOW EXTENDED-RELEASE (ER) TABLETS OR ADMINISTER ER CAPSULES VIA A FEEDING TUBE, OR 2) FAILURE TO ADHERE OR TOLERATE VIA A FEEDING TUBE AN ORAL CARBIDOPA/LEVODOPA REGIMEN. RENEWAL: IMPROVEMENT IN MOTOR SYMPTOMS WHILE ON THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FRUQUINTINIB

---

## Products Affected

- FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# FUTIBATINIB

---

## Products Affected

- LYTGOBI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GALCANEZUMAB-GNLM

## Products Affected

- EMGALITY PEN
- EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.
<b>Other Criteria</b>	MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. EPISODIC CLUSTER HEADACHE: RENEWAL: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GANAXOLONE

---

## Products Affected

- ZTALMY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GEFITINIB

---

## Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# GEPIRONE

## Products Affected

- EXXUA ORAL TABLET EXTENDED  
RELEASE 24 HR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MAJOR DEPRESSIVE DISORDER: INITIAL: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENTS: TRINTELLIX AND ONE GENERIC ANTIDEPRESSANT. INITIAL/RENEWAL: NO CONCURRENT USE WITH ANOTHER 5-HT1A RECEPTOR AGONIST (E.G., BUSPIRONE). RENEWAL: RESPONSE TO OR REMISSION OF DEPRESSIVE SYMPTOMS WITH THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# GILTERITINIB

---

## Products Affected

- XOSPATA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GLASDEGIB

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GLATIRAMER

---

## Products Affected

- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GLP1-DULAGLUTIDE

---

## Products Affected

- TRULICITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GLP1-SEMAGLUTIDE

---

## Products Affected

- OZEMPIC
- RYBELSUS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GLP1-TIRZEPATIDE

---

## Products Affected

- MOUNJARO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# GOSERELIN

## Products Affected

- ZOLADEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS.
<b>Other Criteria</b>	ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# GUSELKUMAB

## Products Affected

- TREMFYA INTRAVENOUS INJECTOR 200 MG/2 ML
- TREMFYA ONE-PRESS • TREMFYA SUBCUTANEOUS SYRINGE
- TREMFYA PEN INDUCTION PK(2PEN)
- TREMFYA PEN SUBCUTANEOUS PEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC), CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

## Products Affected

- *morphine concentrate oral solution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.
<b>Other Criteria</b>	1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# IBRUTINIB

---

## Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CHRONIC GRAFT VS HOST DISEASE (CGVHD): NO CONCURRENT USE WITH JAKAFI, NIKTIMVO, OR REZUROCK.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ICATIBANT

## Products Affected

- *icatibant*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY ANGIOEDEMA (HAE): INITIAL: 1) TYPE III HAE, OR 2) TYPE I OR II HAE CONFIRMED BY ONE OF THE FOLLOWING COMPLEMENT TESTS: C1-INH PROTEIN LEVELS, C4 PROTEIN LEVELS, C1-INH FUNCTIONAL LEVELS, C1Q.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, HEMATOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	HAE: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR THE TREATMENT OF ACUTE HAE ATTACKS. RENEWAL: REDUCTION IN SEVERITY OR DURATION OF ATTACKS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IDELALISIB

---

## Products Affected

- ZYDELIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IMATINIB

---

## Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
Other Criteria	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# IMATINIB SOLUTION

## Products Affected

- IMKELDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
<b>Other Criteria</b>	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR. ALL INDICATIONS: UNABLE TO SWALLOW GENERIC IMATINIB TABLETS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IMETELSTAT

---

## Products Affected

- RYTELO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IMLUNESTRANT

---

## Products Affected

- INLURIYO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# INAVOLISIB

---

## Products Affected

- ITOVEBI ORAL TABLET 3 MG, 9 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# INFLIXIMAB

---

## Products Affected

- *infliximab*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, HUMIRA/CYLTEZO/YUFLYMA, SELARSDI/YESINTEK, XELJANZ, RINVOQ, SKYRIZI, TREMFYA, ORENCIA, OTEZLA. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, HUMIRA/CYLTEZO/YUFLYMA, SELARSDI/YESINTEK, SKYRIZI, TREMFYA, OTEZLA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ. MODERATE TO SEVERE CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: SELARSDI/YESINTEK, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: SELARSDI/YESINTEK, XELJANZ, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA. INITIAL/RENEWAL: RA, PSA, AS, PSO, MODERATE TO SEVERE CD, UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, PSA, AS, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# INSULIN SUPPLIES PAYMENT DETERMINATION

---

## Products Affected

- 1ST TIER UNIFINE PENTP 5MM 31G
- 1ST TIER UNIFINE PNTIP 4MM 32G
- 1ST TIER UNIFINE PNTIP 6MM 31G
- 1ST TIER UNIFINE PNTIP 8MM 31G STRL,SINGLE-USE,SHRT
- 1ST TIER UNIFINE PNTP 29GX1/2"
- 1ST TIER UNIFINE PNTP 31GX3/16
- 1ST TIER UNIFINE PNTP 32GX5/32
- ABOUTTIME PEN NEEDLE
- ADVOCATE INS 0.3 ML 30GX5/16"
- ADVOCATE INS 0.3 ML 31GX5/16"
- ADVOCATE INS 0.5 ML 30GX5/16"
- ADVOCATE INS 0.5 ML 31GX5/16"
- ADVOCATE INS 1 ML 31GX5/16"
- ADVOCATE INS SYR 0.3 ML 29GX1/2
- ADVOCATE INS SYR 0.5 ML 29GX1/2
- ADVOCATE INS SYR 1 ML 29GX1/2"
- ADVOCATE INS SYR 1 ML 30GX5/16
- ADVOCATE PEN NDL 12.7MM 29G
- ADVOCATE PEN NEEDLE 32G 4MM
- ADVOCATE PEN NEEDLE 4MM 33G
- ADVOCATE PEN NEEDLES 5MM 31G
- ADVOCATE PEN NEEDLES 8MM 31G
- ALCOHOL PADS
- ALCOHOL PREP SWABS
- ALCOHOL WIPES
- AQINJECT PEN NEEDLE 31G 5MM
- AQINJECT PEN NEEDLE 32G 4MM
- ASSURE ID DUO PRO NDL 31G 5MM
- ASSURE ID DUO-SHIELD 30GX3/16"
- ASSURE ID DUO-SHIELD 30GX5/16"
- ASSURE ID INSULIN SAFETY SYRINGE 1 ML 29 GAUGE X 1/2"
- ASSURE ID PEN NEEDLE 30GX3/16"
- ASSURE ID PEN NEEDLE 30GX5/16"
- ASSURE ID PEN NEEDLE 31GX3/16"
- ASSURE ID PRO PEN NDL 30G 5MM
- ASSURE ID SYR 0.5 ML 31GX15/64"
- ASSURE ID SYR 1 ML 31GX15/64"
- AUTOSHIELD DUO PEN NDL 30G 5MM
- BD AUTOSHIELD DUO NDL 5MMX30G
- BD ECLIPSE 30GX1/2" SYRINGE
- BD ECLIPSE NEEDLE 30GX1/2" (OTC)
- BD INS SYR 0.3 ML 8MMX31G(1/2)
- BD INS SYR UF 0.3 ML 12.7MMX30G
- BD INS SYR UF 0.5 ML 12.7MMX30G NOT FOR RETAIL SALE
- BD INSULIN SYR 1 ML 27GX12.7MM
- BD INSULIN SYR 1 ML 27GX5/8" MICRO-FINE
- BD LO-DOSE ULTRA-FINE
- BD NANO 2 GEN PEN NDL 32G 4MM
- BD SAFETGLD INS 0.3 ML 29G 13MM
- BD SAFETYGLD INS 0.3 ML 31G 8MM
- BD SAFETYGLD INS 0.5 ML 30G 8MM
- BD SAFETYGLD INS 1 ML 29G 13MM
- BD SAFETYGLID INS 1 ML 6MMX31G
- BD SAFETYGLIDE SYRINGE 27GX5/8
- BD SAFTYGLD INS 0.3 ML 6MMX31G
- BD SAFTYGLD INS 0.5 ML 29G 13MM
- BD SAFTYGLD INS 0.5 ML 6MMX31G
- BD SINGLE USE SWAB
- BD UF MICRO PEN NEEDLE 6MMX32G
- BD UF MINI PEN NEEDLE 5MMX31G
- BD UF NANO PEN NEEDLE 4MMX32G
- BD UF ORIG PEN NDL 12.7MMX29G
- BD UF SHORT PEN NEEDLE 8MMX31G
- BD VEO INS 0.3 ML 6MMX31G (1/2)
- BD VEO INS SYRING 1 ML 6MMX31G
- BD VEO INS SYRN 0.3 ML 6MMX31G
- BD VEO INS SYRN 0.5 ML 6MMX31G
- BORDERED GAUZE 2"X2"
- CAREFINE PEN NEEDLE 12.7MM 29G
- CAREFINE PEN NEEDLE 4MM 32G
- CAREFINE PEN NEEDLE 5MM 32G
- CAREFINE PEN NEEDLE 6MM 31G
- CAREFINE PEN NEEDLE 8MM 30G
- CAREFINE PEN NEEDLES 6MM 32G
- CAREFINE PEN NEEDLES 8MM 31G
- CARETOUCH ALCOHOL 70% PREP PAD
- CARETOUCH PEN NEEDLE 29G 12MM
- CARETOUCH PEN NEEDLE 31GX1/4"

- CARETOUCH PEN NEEDLE 31GX3/16"
- CARETOUCH PEN NEEDLE 31GX5/16"
- CARETOUCH PEN NEEDLE 32GX3/16"
- CARETOUCH PEN NEEDLE 32GX5/32"
- CARETOUCH SYR 0.3 ML 31GX5/16"
- CARETOUCH SYR 0.5 ML 30GX5/16"
- CARETOUCH SYR 0.5 ML 31GX5/16"
- CARETOUCH SYR 1 ML 28GX5/16"
- CARETOUCH SYR 1 ML 29GX5/16"
- CARETOUCH SYR 1 ML 30GX5/16"
- CARETOUCH SYR 1 ML 31GX5/16"
- CLICKFINE PEN NEEDLE 32GX5/32"  
32GX4MM, STERILE
- COMFORT EZ 0.3 ML 31G 15/64"
- COMFORT EZ 0.5 ML 31G 15/64"
- COMFORT EZ INS 0.3 ML 30GX1/2"
- COMFORT EZ INS 0.3 ML 30GX5/16"
- COMFORT EZ INS 1 ML 31G 15/64"
- COMFORT EZ INS 1 ML 31GX5/16"
- COMFORT EZ INSULIN SYR 0.3 ML
- COMFORT EZ INSULIN SYR 0.5 ML
- COMFORT EZ PEN NEEDLE 12MM 29G
- COMFORT EZ PEN NEEDLES 4MM 32G  
SINGLE USE, MICRO
- COMFORT EZ PEN NEEDLES 4MM 33G
- COMFORT EZ PEN NEEDLES 5MM 31G  
MINI
- COMFORT EZ PEN NEEDLES 5MM 32G  
SINGLE USE,MINI,HRI
- COMFORT EZ PEN NEEDLES 5MM 33G
- COMFORT EZ PEN NEEDLES 6MM 31G
- COMFORT EZ PEN NEEDLES 6MM 32G
- COMFORT EZ PEN NEEDLES 6MM 33G
- COMFORT EZ PEN NEEDLES 8MM 31G  
SHORT
- COMFORT EZ PEN NEEDLES 8MM 32G
- COMFORT EZ PEN NEEDLES 8MM 33G
- COMFORT EZ PRO PEN ND L 30G 8MM
- COMFORT EZ PRO PEN ND L 31G 4MM
- COMFORT EZ PRO PEN ND L 31G 5MM
- COMFORT EZ SYR 0.3 ML 29GX1/2"
- COMFORT EZ SYR 0.5 ML 28GX1/2"
- COMFORT EZ SYR 0.5 ML 29GX1/2"
- COMFORT EZ SYR 0.5 ML 30GX1/2"
- COMFORT EZ SYR 1 ML 27G 12.7MM
- COMFORT EZ SYR 1 ML 28GX1/2"
- COMFORT EZ SYR 1 ML 29GX1/2"
- COMFORT EZ SYR 1 ML 30GX1/2"
- COMFORT EZ SYR 1 ML 30GX5/16"
- COMFORT POINT PEN ND L 31GX1/3"
- COMFORT POINT PEN ND L 31GX1/6"
- COMFORT TOUCH PEN ND L 31G 4MM
- COMFORT TOUCH PEN ND L 31G 5MM
- COMFORT TOUCH PEN ND L 31G 6MM
- COMFORT TOUCH PEN ND L 31G 8MM
- COMFORT TOUCH PEN ND L 32G 4MM
- COMFORT TOUCH PEN ND L 32G 5MM
- COMFORT TOUCH PEN ND L 32G 6MM
- COMFORT TOUCH PEN ND L 32G 8MM
- COMFORT TOUCH PEN ND L 33G 4MM
- COMFORT TOUCH PEN ND L 33G 6MM
- COMFORT TOUCH PEN ND L 33GX5MM
- CURAD GAUZE PADS 2" X 2"
- CURITY ALCOHOL PREPS 2  
PLY,MEDIUM
- CURITY GAUZE PADS
- CURITY GAUZE SPONGES (12 PLY)-  
200/BAG
- DERMACEA 2"X2" GAUZE 12 PLY, USP  
TYPE VII
- DERMACEA GAUZE 2"X2" SPONGE 8  
PLY
- DERMACEA NON-WOVEN 2"X2"  
SPNGE
- DROPLET 0.3 ML 29G 12.7MM(1/2)
- DROPLET 0.3 ML 30G 12.7MM(1/2)
- DROPLET 0.5 ML 29GX12.5MM(1/2)
- DROPLET 0.5 ML 30GX12.5MM(1/2)
- DROPLET INS 0.3 ML 29GX12.5MM
- DROPLET INS 0.3 ML 30G 8MM(1/2)
- DROPLET INS 0.3 ML 30GX12.5MM
- DROPLET INS 0.3 ML 31G 6MM(1/2)
- DROPLET INS 0.3 ML 31G 8MM(1/2)
- DROPLET INS 0.5 ML 29G 12.7MM
- DROPLET INS 0.5 ML 30G 12.7MM
- DROPLET INS 0.5 ML 30GX6MM(1/2)
- DROPLET INS 0.5 ML 30GX8MM(1/2)
- DROPLET INS 0.5 ML 31GX6MM(1/2)
- DROPLET INS 0.5 ML 31GX8MM(1/2)
- DROPLET INS SYR 0.3 ML 30GX6MM
- DROPLET INS SYR 0.3 ML 30GX8MM
- DROPLET INS SYR 0.3 ML 31GX6MM
- DROPLET INS SYR 0.3 ML 31GX8MM
- DROPLET INS SYR 0.5 ML 30G 8MM

- DROPLET INS SYR 0.5 ML 31G 6MM
- DROPLET INS SYR 0.5 ML 31G 8MM
- DROPLET INS SYR 1 ML 29G 12.7MM
- DROPLET INS SYR 1 ML 30G 8MM
- DROPLET INS SYR 1 ML 30GX12.5MM
- DROPLET INS SYR 1 ML 30GX6MM
- DROPLET INS SYR 1 ML 31G 6MM
- DROPLET INS SYR 1 ML 31GX8MM
- DROPLET MICRON 34G X 9/64"
- DROPLET PEN NEEDLE 29G 10MM
- DROPLET PEN NEEDLE 29G 12MM
- DROPLET PEN NEEDLE 30G 8MM
- DROPLET PEN NEEDLE 31G 5MM
- DROPLET PEN NEEDLE 31G 6MM
- DROPLET PEN NEEDLE 31G 8MM
- DROPLET PEN NEEDLE 32G 4MM
- DROPLET PEN NEEDLE 32G 5MM
- DROPLET PEN NEEDLE 32G 6MM
- DROPLET PEN NEEDLE 32G 8MM
- DROPSAFE ALCOHOL 70% PREP PADS
- DROPSAFE INS SYR 0.3 ML 31G 6MM
- DROPSAFE INS SYR 0.3 ML 31G 8MM
- DROPSAFE INS SYR 0.5 ML 31G 6MM
- DROPSAFE INS SYR 0.5 ML 31G 8MM
- DROPSAFE INSUL SYR 1 ML 31G 6MM
- DROPSAFE INSUL SYR 1 ML 31G 8MM
- DROPSAFE INSULN 1 ML 29G 12.5MM
- DROPSAFE PEN NEEDLE 31G 5MM
- DROPSAFE PEN NEEDLE 31G 8MM
- DROPSAFE PEN NEEDLE 31GX1/4"
- DRUG MART ULTRA COMFORT SYR
- EASY CMFT SFTY PEN NDL 31G 5MM
- EASY CMFT SFTY PEN NDL 31G 6MM
- EASY CMFT SFTY PEN NDL 32G 4MM
- EASY COMFORT 0.3 ML 31G 1/2"
- EASY COMFORT 0.3 ML 31G 5/16"
- EASY COMFORT 0.3 ML SYRINGE
- EASY COMFORT 0.5 ML 30GX1/2"
- EASY COMFORT 0.5 ML 31GX5/16"
- EASY COMFORT 0.5 ML 32GX5/16"
- EASY COMFORT 0.5 ML SYRINGE
- EASY COMFORT 1 ML 31GX5/16"
- EASY COMFORT 1 ML 32GX5/16"
- EASY COMFORT ALCOHOL 70% PAD
- EASY COMFORT INSULIN 1 ML SYR
- EASY COMFORT PEN NDL 29G 4MM
- EASY COMFORT PEN NDL 29G 5MM
- EASY COMFORT PEN NDL 31GX1/4"
- EASY COMFORT PEN NDL 31GX3/16"
- EASY COMFORT PEN NDL 31GX5/16"
- EASY COMFORT PEN NDL 32GX5/32"
- EASY COMFORT PEN NDL 33G 4MM
- EASY COMFORT PEN NDL 33G 5MM
- EASY COMFORT PEN NDL 33G 6MM
- EASY COMFORT SYR 0.5 ML 29G 8MM
- EASY COMFORT SYR 1 ML 29G 8MM
- EASY COMFORT SYR 1 ML 30GX1/2"
- EASY GLIDE INS 0.3 ML 31GX6MM
- EASY GLIDE INS 0.5 ML 31GX6MM
- EASY GLIDE INS 1 ML 31GX6MM
- EASY GLIDE PEN NEEDLE 4MM 33G
- EASY TOUCH 0.3 ML SYR 30GX1/2"
- EASY TOUCH 0.5 ML SYR 27GX1/2"
- EASY TOUCH 0.5 ML SYR 29GX1/2"
- EASY TOUCH 0.5 ML SYR 30GX1/2"
- EASY TOUCH 0.5 ML SYR 30GX5/16
- EASY TOUCH 1 ML SYR 27GX1/2"
- EASY TOUCH 1 ML SYR 29GX1/2"
- EASY TOUCH 1 ML SYR 30GX1/2"
- EASY TOUCH ALCOHOL 70% PADS  
GAMMA-STERILIZED
- EASY TOUCH FLIPLOK 1 ML 27GX0.5
- EASY TOUCH INSULIN 1 ML 29GX1/2
- EASY TOUCH INSULIN 1 ML 30GX1/2
- EASY TOUCH INSULIN SYR 0.3 ML
- EASY TOUCH INSULIN SYR 0.5 ML
- EASY TOUCH INSULIN SYR 1 ML
- EASY TOUCH INSULIN SYR 1 ML  
RETRACTABLE
- EASY TOUCH INSULN 1 ML 29GX1/2"
- EASY TOUCH INSULN 1 ML 30GX1/2"
- EASY TOUCH INSULN 1 ML 30GX5/16
- EASY TOUCH INSULN 1 ML 31GX5/16
- EASY TOUCH LUER LOK INSUL 1 ML
- EASY TOUCH PEN NEEDLE 29GX1/2"
- EASY TOUCH PEN NEEDLE 30GX5/16
- EASY TOUCH PEN NEEDLE 31GX1/4"
- EASY TOUCH PEN NEEDLE 31GX3/16
- EASY TOUCH PEN NEEDLE 31GX5/16
- EASY TOUCH PEN NEEDLE 32GX1/4"
- EASY TOUCH PEN NEEDLE 32GX3/16
- EASY TOUCH PEN NEEDLE 32GX5/32
- EASY TOUCH SAF PEN NDL 29G 5MM
- EASY TOUCH SAF PEN NDL 29G 8MM

- EASY TOUCH SAF PEN NDL 30G 5MM
- EASY TOUCH SAF PEN NDL 30G 8MM
- EASY TOUCH SYR 0.5 ML 28G 12.7MM
- EASY TOUCH SYR 0.5 ML 29G 12.7MM
- EASY TOUCH SYR 1 ML 27G 16MM
- EASY TOUCH SYR 1 ML 28G 12.7MM
- EASY TOUCH SYR 1 ML 29G 12.7MM
- EASY TOUCH UNI-SLIP SYR 1 ML
- EASYTOUCH SAF PEN NDL 30G 6MM
- EMBRACE PEN NEEDLE 29G 12MM
- EMBRACE PEN NEEDLE 30G 5MM
- EMBRACE PEN NEEDLE 30G 8MM
- EMBRACE PEN NEEDLE 31G 5MM
- EMBRACE PEN NEEDLE 31G 6MM
- EMBRACE PEN NEEDLE 31G 8MM
- EMBRACE PEN NEEDLE 32G 4MM
- EQL INSULIN 0.5 ML SYRINGE
- EQL INSULIN 0.5 ML SYRINGE SHORT NEEDLE
- EXEL U100 0.3 ML 29GX1/2"
- FP INSULIN 1 ML SYRINGE
- FREESTYLE PREC 0.5 ML 30GX5/16
- FREESTYLE PREC 0.5 ML 31GX5/16
- FREESTYLE PREC 1 ML 30GX5/16"
- FREESTYLE PREC 1 ML 31GX5/16"
- GAUZE PAD TOPICAL BANDAGE 2 X 2 "
- GAUZE PADS 2"X2" STRL
- GNP ALCOHOL SWAB STERILE, TWO PLY
- GNP CLICKFINE 31G X 1/4" NDL 6MM, UNIVERSAL
- GNP CLICKFINE 31G X 5/16" NDL 8MM, UNIVERSAL
- GNP PEN NEEDLE 31G 5MM
- GNP PEN NEEDLE 32G 4MM
- GNP PEN NEEDLE 32G 6MM
- GNP SIMPLI PEN NEEDLE 32G 4MM
- GNP ULT C 0.3 ML 29GX1/2" (1/2) 1/2 UNIT
- GNP ULT CMFRT 0.5 ML 29GX1/2"
- GNP ULTRA COMFORT 0.5 ML SYR
- GNP ULTRA COMFORT 1 ML SYRINGE
- GNP ULTRA COMFORT 3/10 ML SYR
- GS PEN NEEDLE 31G X 8MM
- HEALTHWISE INS 0.3 ML 30GX5/16"
- HEALTHWISE INS 0.3 ML 31GX5/16"
- HEALTHWISE INS 0.5 ML 30GX5/16"
- HEALTHWISE INS 0.5 ML 31GX5/16"
- HEALTHWISE INS 1 ML 30GX5/16"
- HEALTHWISE INS 1 ML 31GX5/16"
- HEALTHWISE PEN NEEDLE 31G 5MM
- HEALTHWISE PEN NEEDLE 31G 8MM
- HEALTHWISE PEN NEEDLE 32G 4MM
- HEALTHY ACCENTS PENTIP 4MM 32G
- HEALTHY ACCENTS PENTIP 5MM 31G
- HEALTHY ACCENTS PENTIP 6MM 31G
- HEALTHY ACCENTS PENTIP 8MM 31G
- HEALTHY ACCENTS PENTIP 12MM 29G
- HEB INCONTROL ALCOHOL 70% PADS
- INCONTROL PEN NEEDLE 12MM 29G
- INCONTROL PEN NEEDLE 4MM 32G
- INCONTROL PEN NEEDLE 5MM 31G
- INCONTROL PEN NEEDLE 6MM 31G
- INCONTROL PEN NEEDLE 8MM 31G
- INSULIN 1 ML SYRINGE
- INSULIN SYR 0.3 ML 31GX1/4(1/2)
- INSULIN SYR 0.5 ML 28G 12.7MM (OTC)
- INSULIN SYRIN 0.5 ML 30GX1/2" (RX)
- INSULIN SYRING 0.5 ML 27G 1/2" INNER
- INSULIN SYRINGE 0.3 ML
- INSULIN SYRINGE 0.3 ML 31GX1/4
- INSULIN SYRINGE 0.5 ML
- INSULIN SYRINGE 0.5 ML 31GX1/4
- INSULIN SYRINGE 1 ML
- INSULIN SYRINGE 1 ML 27G 1/2" INNER
- INSULIN SYRINGE 1 ML 27G 16MM
- INSULIN SYRINGE 1 ML 28G 12.7MM (OTC)
- INSULIN SYRINGE 1 ML 30GX1/2" SHORT NEEDLE (OTC)
- INSULIN SYRINGE 1 ML 31GX1/4"
- INSULIN SYRINGE NEEDLELESS
- INSULIN SYRINGE-NEEDLE U-100 SYRINGE 0.3 ML 29 GAUGE, 1 ML 29 GAUGE X 1/2", 1/2 ML 28 GAUGE
- INSULIN U-500 SYRINGE-NEEDLE
- INSUPEN 30G ULTRAFIN NEEDLE
- INSUPEN 31G ULTRAFIN NEEDLE
- INSUPEN 32G 8MM PEN NEEDLE
- INSUPEN PEN NEEDLE 29GX12MM
- INSUPEN PEN NEEDLE 31G 8MM

- INSUPEN PEN NEEDLE 31GX3/16"
- INSUPEN PEN NEEDLE 32G 6MM (RX)
- INSUPEN PEN NEEDLE 32GX4MM
- INSUPEN PEN NEEDLE 33GX4MM
- IV ANTISEPTIC WIPES
- KENDALL ALCOHOL 70% PREP PAD
- LISCO SPONGES 100/BAG
- LITE TOUCH 31GX1/4" PEN NEEDLE
- LITE TOUCH INSULIN 0.5 ML SYR
- LITE TOUCH INSULIN 1 ML SYR
- LITE TOUCH INSULIN SYR 1 ML
- LITE TOUCH PEN NEEDLE 29G
- LITE TOUCH PEN NEEDLE 31G
- LITETOUCH INS 0.3 ML 29GX1/2"
- LITETOUCH INS 0.3 ML 30GX5/16"
- LITETOUCH INS 0.3 ML 31GX5/16"
- LITETOUCH INS 0.5 ML 31GX5/16"
- LITETOUCH SYR 0.5 ML 28GX1/2"
- LITETOUCH SYR 0.5 ML 29GX1/2"
- LITETOUCH SYR 0.5 ML 30GX5/16"
- LITETOUCH SYRIN 1 ML 28GX1/2"
- LITETOUCH SYRIN 1 ML 29GX1/2"
- LITETOUCH SYRIN 1 ML 30GX5/16"
- MAGELLAN INSUL SYRINGE 0.3 ML
- MAGELLAN INSUL SYRINGE 0.5 ML
- MAGELLAN INSULIN SYR 0.3 ML
- MAGELLAN INSULIN SYR 0.5 ML
- MAGELLAN INSULIN SYRINGE 1 ML
- MAXI-COMFORT INS 0.5 ML 28G
- MAXI-COMFORT INS 1 ML 28GX1/2"
- MAXICOMFORT II PEN NDL 31GX6MM
- MAXICOMFORT INS 0.5 ML 27GX1/2"
- MAXICOMFORT INS 1 ML 27GX1/2"
- MAXICOMFORT PEN NDL 29G X 5MM
- MAXICOMFORT PEN NDL 29G X 8MM
- MICRODOT PEN NEEDLE 31GX6MM
- MICRODOT PEN NEEDLE 32GX4MM
- MICRODOT PEN NEEDLE 33GX4MM
- MICRODOT READYGARD NDL 31G 5MM OUTER
- MINI PEN NEEDLE 32G 5MM
- MINI PEN NEEDLE 32G 8MM
- MINI PEN NEEDLE 33G 4MM
- MINI PEN NEEDLE 33G 5MM
- MINI PEN NEEDLE 33G 6MM
- MINI ULTRA-THIN II PEN NDL 31G STERILE
- MONOJECT 0.5 ML SYRN 28GX1/2"
- MONOJECT 1 ML SYRN 27X1/2"
- MONOJECT 1 ML SYRN 28GX1/2" (OTC)
- MONOJECT INSUL SYR U100 (OTC)
- MONOJECT INSUL SYR U100 .5ML,29GX1/2" (OTC)
- MONOJECT INSUL SYR U100 0.5 ML CONVERTS TO 29G (OTC)
- MONOJECT INSUL SYR U100 1 ML
- MONOJECT INSUL SYR U100 1 ML 3'S, 29GX1/2" (OTC)
- MONOJECT INSUL SYR U100 1 ML W/O NEEDLE (OTC)
- MONOJECT INSULIN SYR 0.3 ML
- MONOJECT INSULIN SYR 0.3 ML (OTC)
- MONOJECT INSULIN SYR 0.5 ML
- MONOJECT INSULIN SYR 0.5 ML (OTC)
- MONOJECT INSULIN SYR 1 ML 3'S (OTC)
- MONOJECT INSULIN SYR U-100
- MONOJECT SYRINGE 0.3 ML
- MONOJECT SYRINGE 0.5 ML
- MONOJECT SYRINGE 1 ML
- MS INSULIN SYR 1 ML 31GX5/16" (OTC)
- MS INSULIN SYRINGE 0.3 ML
- NANO 2 GEN PEN NEEDLE 32G 4MM
- NANO PEN NEEDLE 32G 4MM
- NOVOFINE 30
- NOVOFINE 32G NEEDLES
- NOVOFINE PLUS PEN NDL 32GX1/6"
- NOVOTWIST
- PC UNIFINE PENTIPS 8MM NEEDLE SHORT
- PEN NEEDLE 30G 5MM OUTER
- PEN NEEDLE 30G 8MM INNER
- PEN NEEDLE 30G X 5/16"
- PEN NEEDLE 31G X 1/4" HRI
- PEN NEEDLE 6MM 31G 6MM
- PEN NEEDLE, DIABETIC NEEDLE 29 GAUGE X 1/2"
- PEN NEEDLES 12MM 29G 29GX12MM,STRL
- PEN NEEDLES 4MM 32G
- PEN NEEDLES 5MM 31G 31GX5MM,STRL,MINI (OTC)
- PEN NEEDLES 8MM 31G 31GX8MM,STRL,SHORT (OTC)

- PENTIPS PEN NEEDLE 29G 1/2"
- PENTIPS PEN NEEDLE 31G 1/4"
- PENTIPS PEN NEEDLE 31GX3/16" MINI, 5MM
- PENTIPS PEN NEEDLE 31GX5/16" SHORT, 8MM
- PENTIPS PEN NEEDLE 32G 1/4"
- PENTIPS PEN NEEDLE 32GX5/32" 4MM
- PIP PEN NEEDLE 31G X 5MM
- PIP PEN NEEDLE 32G X 4MM
- PREFPLS INS SYR 1 ML 30GX5/16" (OTC)
- PREVENT PEN NEEDLE 31GX1/4"
- PREVENT PEN NEEDLE 31GX5/16"
- PRO COMFORT 0.5 ML 30GX1/2"
- PRO COMFORT 0.5 ML 30GX5/16"
- PRO COMFORT 0.5 ML 31GX5/16"
- PRO COMFORT 1 ML 30GX1/2"
- PRO COMFORT 1 ML 30GX5/16"
- PRO COMFORT 1 ML 31GX5/16"
- PRO COMFORT ALCOHOL 70% PADS
- PRO COMFORT PEN ND L 32G 8MM
- PRO COMFORT PEN ND L 32G X 1/4"
- PRO COMFORT PEN ND L 4MM 32G
- PRO COMFORT PEN ND L 5MM 32G
- PRODIGY INS SYR 1 ML 28GX1/2"
- PRODIGY SYRNG 0.5 ML 31GX5/16"
- PRODIGY SYRNGE 0.3 ML 31GX5/16"
- PURE CMFT SFTY PEN ND L 31G 5MM
- PURE CMFT SFTY PEN ND L 31G 6MM
- PURE CMFT SFTY PEN ND L 32G 4MM
- PURE COMFORT ALCOHOL 70% PADS
- PURE COMFORT PEN ND L 32G 4MM
- PURE COMFORT PEN ND L 32G 5MM
- PURE COMFORT PEN ND L 32G 6MM
- PURE COMFORT PEN ND L 32G 8MM
- RAYA SURE PEN NEEDLE 29G 12MM
- RAYA SURE PEN NEEDLE 31G 4MM
- RAYA SURE PEN NEEDLE 31G 5MM
- RAYA SURE PEN NEEDLE 31G 6MM
- RELI-ON INSULIN 1 ML SYR
- RELION INS SYR 0.3 ML 31GX6MM
- RELION INS SYR 0.5 ML 31GX6MM
- RELION INS SYR 1 ML 31GX15/64"
- SAFESNAP INS SYR UNITS-100 0.3 ML 30GX5/16",10X10
- SAFESNAP INS SYR UNITS-100 0.5 ML 29GX1/2",10X10
- SAFESNAP INS SYR UNITS-100 0.5 ML 30GX5/16",10X10
- SAFESNAP INS SYR UNITS-100 1 ML 28GX1/2",10X10
- SAFESNAP INS SYR UNITS-100 1 ML 29GX1/2",10X10
- SAFETY PEN NEEDLE 31G 4MM
- SAFETY PEN NEEDLE 5MM X 31G
- SAFETY SYRINGE 0.5 ML 30G 1/2"
- SECURES SAFE PEN ND L 30GX5/16" OUTER
- SECURES SAFE SYR 0.5 ML 29G 1/2" OUTER
- SECURES SAFE SYRNG 1 ML 29G 1/2" OUTER
- SKY SAFETY PEN NEEDLE 30G 5MM
- SKY SAFETY PEN NEEDLE 30G 8MM
- SM ULT CFT 0.3 ML 31GX5/16(1/2)
- STERILE PADS 2" X 2"
- SURE CMFT SFTY PEN ND L 31G 6MM
- SURE CMFT SFTY PEN ND L 32G 4MM
- SURE COMFORT 0.5 ML SYRINGE
- SURE COMFORT 1 ML SYRINGE
- SURE COMFORT 3/10 ML SYRINGE
- SURE COMFORT 3/10 ML SYRINGE INSULIN SYRINGE
- SURE COMFORT 30G PEN NEEDLE
- SURE COMFORT ALCOHOL PREP PADS
- SURE COMFORT INS 0.3 ML 31GX1/4
- SURE COMFORT INS 0.5 ML 31GX1/4
- SURE COMFORT INS 1 ML 31GX1/4"
- SURE COMFORT PEN ND L 29GX1/2" 12.7MM
- SURE COMFORT PEN ND L 31G 5MM
- SURE COMFORT PEN ND L 31G 8MM
- SURE COMFORT PEN ND L 32G 4MM
- SURE COMFORT PEN ND L 32G 6MM
- SURE-FINE PEN NEEDLES 12.7MM
- SURE-FINE PEN NEEDLES 5MM
- SURE-FINE PEN NEEDLES 8MM
- SURE-JECT INSU SYR U100 0.3 ML
- SURE-JECT INSU SYR U100 0.5 ML
- SURE-JECT INSU SYR U100 1 ML
- SURE-JECT INSUL SYR U100 1 ML
- SURE-JECT INSULIN SYRINGE 1 ML
- SURE-PREP ALCOHOL PREP PADS

- TECHLITE 0.3 ML 29GX12MM (1/2)
- TECHLITE 0.3 ML 30GX8MM (1/2)
- TECHLITE 0.3 ML 31GX6MM (1/2)
- TECHLITE 0.3 ML 31GX8MM (1/2)
- TECHLITE 0.5 ML 30GX12MM (1/2)
- TECHLITE 0.5 ML 30GX8MM (1/2)
- TECHLITE 0.5 ML 31GX6MM (1/2)
- TECHLITE 0.5 ML 31GX8MM (1/2)
- TECHLITE INS SYR 1 ML 29GX12MM
- TECHLITE INS SYR 1 ML 30GX12MM
- TECHLITE INS SYR 1 ML 31GX6MM
- TECHLITE INS SYR 1 ML 31GX8MM
- TECHLITE PEN NEEDLE 29GX1/2"
- TECHLITE PEN NEEDLE 29GX3/8"
- TECHLITE PEN NEEDLE 31GX1/4"
- TECHLITE PEN NEEDLE 31GX3/16"
- TECHLITE PEN NEEDLE 31GX5/16"
- TECHLITE PEN NEEDLE 32GX1/4"
- TECHLITE PEN NEEDLE 32GX5/16"
- TECHLITE PEN NEEDLE 32GX5/32"
- TECHLITE PLUS PEN NDL 32G 4MM
- TERUMO INS SYRINGE U100-1 ML
- TERUMO INS SYRINGE U100-1/2 ML
- TERUMO INS SYRINGE U100-1/3 ML
- TERUMO INS SYRNG U100-1/2 ML
- THINPRO INS SYRIN U100-0.3 ML
- THINPRO INS SYRIN U100-0.5 ML
- THINPRO INS SYRIN U100-1 ML
- TOPCARE CLICKFINE 31G X 1/4"
- TOPCARE CLICKFINE 31G X 5/16"
- TOPCARE ULTRA COMFORT SYRINGE
- TRUE CMFRT PRO 0.5 ML 30G 5/16"
- TRUE CMFRT PRO 0.5 ML 31G 5/16"
- TRUE CMFRT PRO 0.5 ML 32G 5/16"
- TRUE CMFT SFTY PEN NDL 31G 5MM
- TRUE CMFT SFTY PEN NDL 31G 6MM
- TRUE CMFT SFTY PEN NDL 32G 4MM
- TRUE COMFORT 0.5 ML 30G 1/2"
- TRUE COMFORT 0.5 ML 30G 5/16"
- TRUE COMFORT 0.5 ML 31G 5/16"
- TRUE COMFORT 0.5 ML 31GX5/16"
- TRUE COMFORT 1 ML 31GX5/16"
- TRUE COMFORT ALCOHOL 70% PADS
- TRUE COMFORT PEN NDL 31G 8MM
- TRUE COMFORT PEN NDL 31GX5MM
- TRUE COMFORT PEN NDL 31GX6MM
- TRUE COMFORT PEN NDL 32G 5MM
- TRUE COMFORT PEN NDL 32G 6MM
- TRUE COMFORT PEN NDL 32GX4MM
- TRUE COMFORT PEN NDL 33G 4MM
- TRUE COMFORT PEN NDL 33G 5MM
- TRUE COMFORT PEN NDL 33G 6MM
- TRUE COMFORT PRO 1 ML 30G 1/2"
- TRUE COMFORT PRO 1 ML 30G 5/16"
- TRUE COMFORT PRO 1 ML 31G 5/16"
- TRUE COMFORT PRO 1 ML 32G 5/16"
- TRUE COMFORT PRO ALCOHOL PADS
- TRUE COMFORT SFTY 1 ML 30G 1/2"
- TRUE COMFRT PRO 0.5 ML 30G 1/2"
- TRUE COMFRT SFTY 1 ML 30G 5/16"
- TRUE COMFRT SFTY 1 ML 31G 5/16"
- TRUE COMFRT SFTY 1 ML 32G 5/16"
- TRUEPLUS PEN NEEDLE 29GX1/2"
- TRUEPLUS PEN NEEDLE 31G X 1/4"
- TRUEPLUS PEN NEEDLE 31GX3/16"
- TRUEPLUS PEN NEEDLE 31GX5/16"
- TRUEPLUS PEN NEEDLE 32GX5/32"
- TRUEPLUS SYR 0.3 ML 29GX1/2"
- TRUEPLUS SYR 0.3 ML 30GX5/16"
- TRUEPLUS SYR 0.3 ML 31GX5/16"
- TRUEPLUS SYR 0.5 ML 28GX1/2"
- TRUEPLUS SYR 0.5 ML 29GX1/2"
- TRUEPLUS SYR 0.5 ML 30GX5/16"
- TRUEPLUS SYR 0.5 ML 31GX5/16"
- TRUEPLUS SYR 1 ML 28GX1/2"
- TRUEPLUS SYR 1 ML 29GX1/2"
- TRUEPLUS SYR 1 ML 30GX5/16"
- TRUEPLUS SYR 1 ML 31GX5/16"
- ULTICAR INS 0.3 ML 31GX1/4(1/2)
- ULTICARE INS 1 ML 31GX1/4"
- ULTICARE INS SYR 0.3 ML 30G 8MM
- ULTICARE INS SYR 0.3 ML 31G 6MM
- ULTICARE INS SYR 0.3 ML 31G 8MM
- ULTICARE INS SYR 0.5 ML 30G 8MM (OTC)
- ULTICARE INS SYR 0.5 ML 31G 6MM
- ULTICARE INS SYR 0.5 ML 31G 8MM (OTC)
- ULTICARE INS SYR 1 ML 30GX1/2"
- ULTICARE PEN NEEDLE 31GX3/16"
- ULTICARE PEN NEEDLE 6MM 31G
- ULTICARE PEN NEEDLE 8MM 31G
- ULTICARE PEN NEEDLES 12MM 29G
- ULTICARE PEN NEEDLES 4MM 32G

- MICRO, 32GX4MM
- ULTICARE PEN NEEDLES 6MM 32G
  - ULTICARE SAFE PEN NDL 30G 8MM
  - ULTICARE SAFE PEN NDL 5MM 30G
  - ULTICARE SAFETY 0.5 ML 29GX1/2 (RX)
  - ULTICARE SYR 0.3 ML 29G 12.7MM
  - ULTICARE SYR 0.3 ML 30GX1/2"
  - ULTICARE SYR 0.3 ML 31GX5/16" SHORT NDL
  - ULTICARE SYR 0.5 ML 30GX1/2"
  - ULTICARE SYR 0.5 ML 31GX5/16" SHORT NDL
  - ULTICARE SYR 1 ML 31GX5/16"
  - ULTIGUARD SAFE 1 ML 30G 12.7MM
  - ULTIGUARD SAFE0.3 ML 30G 12.7MM
  - ULTIGUARD SAFE0.5 ML 30G 12.7MM
  - ULTIGUARD SAFEPACK 1 ML 31G 8MM
  - ULTIGUARD SAFEPACK 29G 12.7MM
  - ULTIGUARD SAFEPACK 31G 5MM
  - ULTIGUARD SAFEPACK 31G 6MM
  - ULTIGUARD SAFEPACK 31G 8MM
  - ULTIGUARD SAFEPACK 32G 4MM
  - ULTIGUARD SAFEPACK 32G 6MM
  - ULTIGUARD SAFEPK 0.3 ML 31G 8MM
  - ULTIGUARD SAFEPK 0.5 ML 31G 8MM
  - ULTILET ALCOHOL STERL SWAB
  - ULTILET INSULIN SYRINGE 0.3 ML
  - ULTILET INSULIN SYRINGE 0.5 ML
  - ULTILET INSULIN SYRINGE 1 ML
  - ULTILET PEN NEEDLE
  - ULTILET PEN NEEDLE 4MM 32G
  - ULTRA COMFORT 0.3 ML SYRINGE
  - ULTRA COMFORT 0.5 ML 28GX1/2" CONVERTS TO 29G
  - ULTRA COMFORT 0.5 ML 29GX1/2"
  - ULTRA COMFORT 0.5 ML SYRINGE
  - ULTRA COMFORT 1 ML 31GX5/16"
  - ULTRA COMFORT 1 ML SYRINGE
  - ULTRA FLO 0.3 ML 30G 1/2" (1/2)
  - ULTRA FLO 0.3 ML 30G 5/16"(1/2)
  - ULTRA FLO 0.3 ML 31G 5/16"(1/2)
  - ULTRA FLO PEN NEEDLE 31G 5MM
  - ULTRA FLO PEN NEEDLE 31G 8MM
  - ULTRA FLO PEN NEEDLE 32G 4MM
  - ULTRA FLO PEN NEEDLE 33G 4MM
  - ULTRA FLO PEN NEEDLES 12MM 29G
  - ULTRA FLO SYR 0.3 ML 29GX1/2"
  - ULTRA FLO SYR 0.3 ML 30G 5/16"
  - ULTRA FLO SYR 0.3 ML 31G 5/16"
  - ULTRA FLO SYR 0.5 ML 29G 1/2"
  - ULTRA THIN PEN NDL 32G X 4MM
  - ULTRA-FINE 0.3 ML 30G 12.7MM
  - ULTRA-FINE 0.3 ML 31G 6MM (1/2)
  - ULTRA-FINE 0.3 ML 31G 8MM (1/2)
  - ULTRA-FINE 0.5 ML 30G 12.7MM
  - ULTRA-FINE INS SYR 1 ML 31G 6MM
  - ULTRA-FINE INS SYR 1 ML 31G 8MM
  - ULTRA-FINE PEN NDL 29G 12.7MM
  - ULTRA-FINE PEN NEEDLE 31G 5MM
  - ULTRA-FINE PEN NEEDLE 31G 8MM
  - ULTRA-FINE PEN NEEDLE 32G 6MM
  - ULTRA-FINE SYR 0.5 ML 31G 6MM
  - ULTRA-FINE SYR 0.5 ML 31G 8MM
  - ULTRA-FINE SYR 1 ML 30G 12.7MM
  - ULTRA-THIN II 1 ML 31GX5/16"
  - ULTRA-THIN II INS 0.3 ML 30G
  - ULTRA-THIN II INS 0.3 ML 31G
  - ULTRA-THIN II INS 0.5 ML 29G
  - ULTRA-THIN II INS 0.5 ML 30G
  - ULTRA-THIN II INS 0.5 ML 31G
  - ULTRA-THIN II INS SYR 1 ML 29G
  - ULTRA-THIN II INS SYR 1 ML 30G
  - ULTRA-THIN II PEN NDL 29GX1/2"
  - ULTRA-THIN II PEN NDL 31GX5/16"
  - ULTRACARE INS 0.3 ML 30GX5/16"
  - ULTRACARE INS 0.3 ML 31GX5/16"
  - ULTRACARE INS 0.5 ML 30GX1/2"
  - ULTRACARE INS 0.5 ML 30GX5/16"
  - ULTRACARE INS 0.5 ML 31GX5/16"
  - ULTRACARE INS 1 ML 30G X 5/16"
  - ULTRACARE INS 1 ML 30GX1/2"
  - ULTRACARE INS 1 ML 31G X 5/16"
  - ULTRACARE PEN NEEDLE 31GX1/4"
  - ULTRACARE PEN NEEDLE 31GX3/16"
  - ULTRACARE PEN NEEDLE 31GX5/16"
  - ULTRACARE PEN NEEDLE 32GX1/4"
  - ULTRACARE PEN NEEDLE 32GX3/16"
  - ULTRACARE PEN NEEDLE 32GX5/32"
  - ULTRACARE PEN NEEDLE 33GX5/32"
  - UNIFINE OTC PEN NEEDLE 31G 5MM
  - UNIFINE OTC PEN NEEDLE 32G 4MM
  - UNIFINE PEN NEEDLE 32G 4MM
  - UNIFINE PENTIPS 12MM 29G

- 29GX12MM, STRL
- UNIFINE PENTIPS 31GX3/16" 31GX5MM,STRL,MINI
- UNIFINE PENTIPS 32G 4MM
- UNIFINE PENTIPS 32GX1/4"
- UNIFINE PENTIPS 33GX5/32"
- UNIFINE PENTIPS 6MM 31G
- UNIFINE PENTIPS MAX 30GX3/16"
- UNIFINE PENTIPS NEEDLES 29G 12MM
- UNIFINE PENTIPS PLUS 29GX1/2" 12MM
- UNIFINE PENTIPS PLUS 30GX3/16"
- UNIFINE PENTIPS PLUS 31GX1/4" ULTRA SHORT, 6MM
- UNIFINE PENTIPS PLUS 31GX3/16" MINI
- UNIFINE PENTIPS PLUS 31GX5/16" SHORT
- UNIFINE PENTIPS PLUS 32GX5/32"
- UNIFINE PENTIPS PLUS 33GX5/32"
- UNIFINE PROTECT 30G 5MM
- UNIFINE PROTECT 30G 8MM
- UNIFINE PROTECT 32G 4MM
- UNIFINE SAFECONTROL 30G 5MM
- UNIFINE SAFECONTROL 30G 8MM
- UNIFINE SAFECONTROL 31G 5MM
- UNIFINE SAFECONTROL 31G 6MM
- UNIFINE SAFECONTROL 31G 8MM
- UNIFINE SAFECONTROL 32G 4MM
- UNIFINE ULTRA PEN NDL 31G 5MM
- UNIFINE ULTRA PEN NDL 31G 6MM
- UNIFINE ULTRA PEN NDL 31G 8MM
- UNIFINE ULTRA PEN NDL 32G 4MM
- VANISHPOINT 0.5 ML 30GX1/2" SY OUTER
- VANISHPOINT INS 1 ML 30GX3/16"
- VANISHPOINT U-100 29X1/2 SYR
- VERIFINE INS SYR 1 ML 29G 1/2"
- VERIFINE PEN NEEDLE 29G 12MM
- VERIFINE PEN NEEDLE 31G 5MM
- VERIFINE PEN NEEDLE 31G X 6MM
- VERIFINE PEN NEEDLE 31G X 8MM
- VERIFINE PEN NEEDLE 32G 6MM
- VERIFINE PEN NEEDLE 32G X 4MM
- VERIFINE PEN NEEDLE 32G X 5MM
- VERIFINE PLUS PEN NDL 31G 5MM
- VERIFINE PLUS PEN NDL 31G 8MM
- VERIFINE PLUS PEN NDL 32G 4MM
- VERIFINE PLUS PEN NDL 32G 4MM-SHARPS CONTAINER
- VERIFINE SYRING 0.5 ML 29G 1/2"
- VERIFINE SYRING 1 ML 31G 5/16"
- VERIFINE SYRNG 0.3 ML 31G 5/16"
- VERIFINE SYRNG 0.5 ML 31G 5/16"
- VERSALON ALL PURPOSE SPONGE 25'S,N-STERILE,3PLY
- WEBCOL ALCOHOL PREPS 20'S,LARGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	ONLY COVERED UNDER PART D WHEN USED CONCURRENTLY WITH INSULIN.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	LIFETIME

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# INTERFERON FOR MS-AVONEX

---

## Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# INTERFERON FOR MS-BETASERON

---

## Products Affected

- BETASERON SUBCUTANEOUS KIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# INTERFERON FOR MS-PLEGRIDY

---

## Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# INTERFERON GAMMA-1B

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST OR HEMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IPILIMUMAB

---

## Products Affected

- YERVOY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO
Other Criteria	RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ISAVUCONAZONIUM

---

## Products Affected

- CRESEMBA ORAL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INVASIVE ASPERGILLOSIS, INVASIVE MUCORMYCOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	6 MONTHS
<b>Other Criteria</b>	INVASIVE ASPERGILLOSIS: TRIAL OF OR CONTRAINDICATION TO VORICONAZOLE. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# IVACAFTOR

---

## Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CYSTIC FIBROSIS (CF): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: INITIAL: 1) NOT HOMOZYGOUS FOR F508DEL MUTATION IN THE CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR (CFTR) GENE, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR. RENEWAL: 1) IMPROVEMENT IN CLINICAL STATUS, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# IVOSIDENIB

---

## Products Affected

- TIBSOVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IXAZOMIB

---

## Products Affected

- NINLARO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LAMOTRIGINE

---

## Products Affected

- SUBVENITE ORAL SUSPENSION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ALL INDICATIONS: CONTRAINDICATION TO OR UNABLE TO SWALLOW LAMOTRIGINE TABLETS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LANREOTIDE

## Products Affected

- *lanreotide subcutaneous syringe 120 mg/0.5 ml*
- SOMATULINE DEPOT SUBCUTANEOUS SYRINGE 60 MG/0.2 ML, 90 MG/0.3 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ACROMEGALY: INITIAL/RENEWAL: 12 MOS. GEP-NETS, CARCINOID SYNDROME: 12 MOS.
Other Criteria	ACROMEGALY: RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# LAPATINIB

---

## Products Affected

- *lapatinib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LAROTRECTINIB

---

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	VITRAKVI ORAL SOLUTION: 1) TRIAL OF VITRAKVI CAPSULES, OR 2) UNABLE TO TAKE CAPSULE FORMULATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LAZERTINIB

---

## Products Affected

- LAZCLUZE ORAL TABLET 240 MG, 80 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LEDIPASVIR-SOFOSBUVIR

## Products Affected

- HARVONI ORAL PELLETS IN PACKET  
33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANAVIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LENALIDOMIDE

---

## Products Affected

- *lenalidomide*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LENVATINIB

---

## Products Affected

- LENVIMA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LETERMOVIR

## Products Affected

- PREVYMIS ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	HSCT: NOT AT RISK FOR LATE CMV: 4 MOS, AT RISK FOR LATE CMV: 7 MOS. KIDNEY TRANSPLANT: 7 MOS.
<b>Other Criteria</b>	HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 100 DAYS POST TRANSPLANT IF NOT AT RISK FOR LATE CYTOMEGALOVIRUS (CMV) INFECTION AND DISEASE, OR BEYOND 200 DAYS POST TRANSPLANT IF AT RISK FOR LATE CMV INFECTION AND DISEASE. KIDNEY TRANSPLANT: 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 7 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 200 DAYS POST TRANSPLANT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LEUPROLIDE

---

## Products Affected

- *leuprolide subcutaneous kit*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	PROSTATE CANCER: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LEUPROLIDE DEPOT

---

## Products Affected

- *leuprolide acetate (3 month)*
- LUTRATE DEPOT (3 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LEUPROLIDE MESYLATE

---

## Products Affected

- CAMCEVI (6 MONTH)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LEUPROLIDE-ELIGARD

---

## Products Affected

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LEUPROLIDE-LUPRON DEPOT

## Products Affected

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.
<b>Other Criteria</b>	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# LEUPROLIDE-LUPRON DEPOT-PED

## Products Affected

- LUPRON DEPOT-PED (3 MONTH)
- LUPRON DEPOT-PED  
INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVEL OF LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVEL OF LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# L-GLUTAMINE

## Products Affected

- *glutamine (sickle cell)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
<b>Coverage Duration</b>	INITIAL: 12 MONTHS. RENEWAL: LIFETIME.
<b>Other Criteria</b>	SCD: INITIAL: AGES 18 YEARS OR OLDER: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. AGES 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SCD.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LIDOCAINE OINTMENT

---

## Products Affected

- *lidocaine topical ointment*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LIDOCAINE PATCH

## Products Affected

- *dermacinrx lidocan 5% patch outer*
- *lidocaine topical adhesive patch,medicated 5 %*
- *lidocan iii*
- *tridacaine ii*
- ZTLIDO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	1) PAIN ASSOCIATED WITH POST-HERPETIC NEURALGIA, 2) NEUROPATHY DUE TO DIABETES MELLITUS, 3) CHRONIC BACK PAIN, OR 4) OSTEOARTHRITIS OF THE KNEE OR HIP.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LIDOCAINE PRILOCAINE

---

## Products Affected

- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LINVOSELTAMAB-GCPT

---

## Products Affected

- LYNOZYFIC INTRAVENOUS SOLUTION 2 MG/ML, 20 MG/ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LONCASTUXIMAB TESIRINE-LPYL

---

## Products Affected

- ZYNLONTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LORLATINIB

---

## Products Affected

- LORBRENA ORAL TABLET 100 MG, 25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LOTILANER

---

## Products Affected

- XDEMZY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	DEMODEX BLEPHARITIS: 18 YEARS OF AGE OR OLDER
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 WEEKS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LUMACAFITOR-IVACAFITOR

## Products Affected

- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CYSTIC FIBROSIS (CF): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
<b>Other Criteria</b>	CF: INITIAL: NO CONCURRENT USE WITH ANOTHER CYSTIC FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR (CFTR) MODULATOR. RENEWAL: 1) IMPROVEMENT IN CLINICAL STATUS, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MACITENTAN

## Products Affected

- OPSUMIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MARGETUXIMAB-CMKB

---

## Products Affected

- MARGENZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MARIBAVIR

---

## Products Affected

- LIVTENCITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MAVACAMTEN

## Products Affected

- CAMZYOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	OBSTRUCTIVE HYPERTROPHIC CARDIOMYOPATHY(HCM): INITIAL: LEFT VENTRICULAR OUTFLOW TRACK (LVOT) GRADIENT OF 50 MMHG OR HIGHER
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	OBSTRUCTIVE HCM: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	OBSTRUCTIVE HCM: INITIAL: TRIAL OF, CONTRAINDICATION, OR INTOLERANCE TO A BETA-BLOCKER OR A NON-DIHYDROPYRIDINE CALCIUM CHANNEL BLOCKER. RENEWAL: CONTINUED CLINICAL BENEFIT (E.G., REDUCTION OF SYMPTOMS, NYHA CLASSIFICATION IMPROVEMENT).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# MECASERMIN

## Products Affected

- INCRELEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	GROWTH FAILURE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST OR NEPHROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	GROWTH FAILURE: INITIAL: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF WRIST AND HAND. INITIAL/RENEWAL: NO CONCURRENT USE WITH ANOTHER GROWTH HORMONE MEDICATION. RENEWAL: IMPROVEMENT WHILE ON THERAPY (INCREASE IN HEIGHT OR INCREASE IN HEIGHT VELOCITY).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MECHLORETHAMINE

---

## Products Affected

- VALCHLOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MEPOLIZUMAB

## Products Affected

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOSINOPHILIC COPD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL: CRSWNP: 6 MO. OTHERS: 12 MO. RENEWAL: CRSWNP, ASTHMA, COPD, EGPA, HES: 12 MO.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING AT LEAST 3 DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA. CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP): 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, 2) EVIDENCE OF NASAL POLYPS BY DIRECT EXAMINATION, ENDOSCOPY, OR SINUS CT SCAN, AND 3) INADEQUATELY CONTROLLED DISEASE. EOSINOPHILIC COPD: USED IN COMBINATION WITH A LAMA/LABA/ICS. RENEWAL: ASTHMA: 1) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, AND 3) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION. CRSWNP: 1) CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH ANOTHER</p>

PA Criteria	Criteria Details
	<p>SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS (EGPA): 1) REDUCTION IN EGPA SYMPTOMS COMPARED TO BASELINE OR ABILITY TO REDUCE/ELIMINATE CORTICOSTEROID USE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION. EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A LAMA/LABA/ICS, 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY (A) REDUCTION IN COPD EXACERBATIONS FROM BASELINE, (B) REDUCTION IN SEVERITY OR FREQUENCY OF COPD-RELATED SYMPTOMS, OR (C) INCREASE IN FEV1 OF AT LEAST 5 PERCENT FROM PRETREATMENT BASELINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# METYROSINE

## Products Affected

- *metyrosine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PHEOCHROMOCYTOMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, ENDOCRINE SURGEON, OR HEMATOLOGIST-ONCOLOGIST.
<b>Coverage Duration</b>	PREOPERATIVE PREPARATION FOR SURGERY: 30 DAYS. MALIGNANT PHEOCHROMOCYTOMA: INITIAL/RENEWAL:12 MOS.
<b>Other Criteria</b>	PHEOCHROMOCYTOMA: INITIAL: HAS NON-METASTATIC PHEOCHROMOCYTOMA. PREOPERATIVE PREPARATION FOR SURGERY: USE IN COMBINATION WITH AN ALPHA-ADRENERGIC RECEPTOR BLOCKER. RENEWAL: MALIGNANT PHEOCHROMOCYTOMA: STABLE OR CLINICAL IMPROVEMENT WHILE ON THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MIDOSTAURIN

---

## Products Affected

- RYDAPT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MIFEPRISTONE

## Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY: 1) 24-HR URINE FREE CORTISOL (AT LEAST 2 TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (AT LEAST 2 TESTS TO CONFIRM).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE), 2) CONTINUES TO HAVE TOLERABILITY TO THERAPY, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MILTEFOSINE

---

## Products Affected

- IMPAVIDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MIRDAMETINIB

---

## Products Affected

- GOMEKLI ORAL CAPSULE 1 MG, 2 MG
- GOMEKLI ORAL TABLET FOR SUSPENSION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MIRVETUXIMAB SORAVTANSINE-GYNX

## Products Affected

- ELAHERE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: AN OPHTHALMIC EXAM, INCLUDING VISUAL ACUITY AND SLIT LAMP EXAM, WILL BE COMPLETED PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MOMELOTINIB

---

## Products Affected

- OJAARA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# MOSUNETUZUMAB-AXGB

## Products Affected

- LUNSUMIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS.
<b>Other Criteria</b>	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NARCOLEPSY AGENTS

---

## Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NAXITAMAB-GQGK

---

## Products Affected

- DANYELZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NERATINIB

---

## Products Affected

- NERLYNX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# NILOTINIB - TASIGNA

---

## Products Affected

- TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND MEDICATION IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NILOTINIB-DANZITEN

## Products Affected

- DANZITEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): 1) PERFORMED MUTATIONAL ANALYSIS PRIOR TO INITIATION OF THERAPY, AND 2) THERAPY IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NINTEDANIB

## Products Affected

- OFEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): AT LEAST 10% FIBROSIS ON A CHEST HRCT.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG-INDUCED LUNG TOXICITY, RECURRENT ASPIRATION). PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENED/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# NIRAPARIB

---

## Products Affected

- ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# NIRAPARIB-ABIRATERONE

## Products Affected

- AKEEGA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC), METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NIROGACESTAT

---

## Products Affected

- OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NITISINONE

## Products Affected

- *nitisinone*
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# NIVOLUMAB

---

## Products Affected

- OPDIVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NIVOLUMAB-HYALURONIDASE-NVHY

---

## Products Affected

- OPDIVO QVANTIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NIVOLUMAB-RELATLIMAB-RMBW

---

## Products Affected

- OPDUALAG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NOGAPENDEKIN ALFA

---

## Products Affected

- ANKTIVA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	40 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OFATUMUMAB SQ

---

## Products Affected

- KESIMPTA PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OLAPARIB

## Products Affected

- LYNPARZA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: MEDICATION WILL BE USED AS MONOTHERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OLUTASIDENIB

---

## Products Affected

- REZLIDHIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OMACETAXINE

---

## Products Affected

- SYNRIBO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OMALIZUMAB

## Products Affected

- XOLAIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL OF AT LEAST 30 IU/ML. FOOD ALLERGY: 1) IGE SERUM LEVEL OF AT LEAST 30 IU/ML, AND 2) POSITIVE SKIN PRICK TEST TO AT LEAST ONE FOOD, OR POSITIVE MEDICALLY MONITORED FOOD CHALLENGE TO AT LEAST ONE FOOD.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL/RENEWAL: CHRONIC SPONTANEOUS URTICARIA (CSU): PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, DERMATOLOGIST, OR IMMUNOLOGIST. INITIAL: CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP): PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. FOOD ALLERGY: PRESCRIBED BY OR IN CONSULTATION WITH ALLERGIST OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: ASTHMA 12 MO/12 MO, CSU 6 MO/12 MO, CRSWNP 6 MO/12 MO, FOOD ALLERGY 12 MO/24 MO

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: CSU: 1) TRIAL OF AND MAINTAINED ON, OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE, 2) STILL EXPERIENCES HIVES OR ANGIOEDEMA ON MOST DAYS OF THE WEEK FOR AT LEAST 6 WEEKS, AND 3) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: DUPIXENT. CRSWNP: 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: NUCALA, DUPIXENT, 3) EVIDENCE OF NASAL POLYPS BY DIRECT EXAMINATION, ENDOSCOPY, OR SINUS CT SCAN, AND 4) INADEQUATELY CONTROLLED DISEASE. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING AT LEAST 3 DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA. FOOD ALLERGY: CONCURRENT USE WITH AN ACTIVE PRESCRIPTION FOR EPINEPHRINE AUTO-INJECTOR/INJECTION .</p> <p>INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: CSU: MAINTAINED ON OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE. CRSWNP: CLINICAL BENEFIT COMPARED TO BASELINE. ASTHMA: 1) CONTINUED USE OF ICS AND AT LEAST</p>

PA Criteria	Criteria Details
	ONE OTHER MAINTENANCE MEDICATION, AND 2) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE. FOOD ALLERGY: 1) PERSISTENT IGE-MEDIATED FOOD ALLERGY, AND 2) CONCURRENT USE WITH AN ACTIVE PRESCRIPTION FOR EPINEPHRINE AUTO-INJECTOR/INJECTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# OSIMERTINIB

---

## Products Affected

- TAGRISSO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OXANDROLONE

## Products Affected

- *oxandrolone*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 MONTHS
<b>Other Criteria</b>	PROTEIN CATABOLISM, BONE PAIN: 1) MONITORED FOR PELIOSIS HEPATIS, LIVER CELL TUMORS, AND BLOOD LIPID CHANGES, 2) DOES NOT HAVE KNOWN OR SUSPECTED: CARCINOMA OF THE PROSTATE OR BREAST IN MALE PATIENTS, CARCINOMA OF THE BREAST IN FEMALES WITH HYPERCALCEMIA, NEPHROSIS (THE NEPHROTIC PHASE OF NEPHRITIS), OR HYPERCALCEMIA, AND 3) DOES NOT HAVE SEVERE HEPATIC DYSFUNCTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PACRITINIB

---

## Products Affected

- VONJO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PALBOCICLIB

---

## Products Affected

- IBRANCE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PARATHYROID HORMONE

## Products Affected

- NATPARA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PASIREOTIDE DIASPARTATE

---

## Products Affected

- SIGNIFOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PAZOPANIB

---

## Products Affected

- *pazopanib oral tablet 200 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PEGFILGRASTIM - APGF

---

## Products Affected

- NYVEPRIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PEGFILGRASTIM - CBQV

## Products Affected

- UDENYCA ONBODY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PREScribed BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: UDENYCA: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NYVEPRIA. UDENYCA ONBODY: 1) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NYVEPRIA, OR 2) BARRIER TO ACCESS (E.G., TRAVEL BARRIERS, UNABLE TO RETURN TO CLINIC FOR INJECTIONS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PEGINTERFERON ALFA-2A

---

## Products Affected

- PEGASYS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HEPATITIS B: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, OR PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G., HEPATOLOGIST).
<b>Coverage Duration</b>	HEP B/HEP C: 48 WEEKS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PEGVISOMANT

---

## Products Affected

- SOMAVERT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PEMBROLIZUMAB

---

## Products Affected

- KEYTRUDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PEMBROLIZUMAB-BERAHYALURONIDASE ALFA-PMPH

---

## Products Affected

- KEYTRUDA QLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PEMIGATINIB

---

## Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CHOLANGIOCARCINOMA, MYELOID/LYMPHOID NEOPLASMS: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PENICILLAMINE TABLET

## Products Affected

- *penicillamine oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CYSTINURIA: HAS NEPHROLITHIASIS AND ONE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTINE, 2) PRESENCE OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) FAMILY HISTORY OF CYSTINURIA AND POSITIVE CYANIDE-NITROPRUSSIDE SCREENING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
<b>Other Criteria</b>	INITIAL: WILSONS DISEASE: 1) LEIPZIG SCORE OF 4 OR GREATER. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: CONTINUES TO BENEFIT FROM THE MEDICATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PEXIDARTINIB

---

## Products Affected

- TURALIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PIMAVANSERIN

---

## Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER
Prescriber Restrictions	PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST).
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PIRFENIDONE

## Products Affected

- *pirfenidone oral capsule*
- *pirfenidone oral tablet 267 mg, 534 mg, 801 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# PIRTOBRUTINIB

---

## Products Affected

- JAYPIRCA ORAL TABLET 100 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# POMALIDOMIDE

---

## Products Affected

- POMALYST

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PONATINIB

---

## Products Affected

- ICLUSIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CHRONIC MYELOID LEUKEMIA (CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# POSACONAZOLE TABLET

---

## Products Affected

- *posaconazole oral tablet, delayed release (dr/ec)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE, PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PRALSETINIB

---

## Products Affected

- GAVRETO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PYRIMETHAMINE

## Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.
<b>Other Criteria</b>	TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# QUININE

---

## Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# QUIZARTINIB

---

## Products Affected

- VANFLYTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# REGORAFENIB

---

## Products Affected

- STIVARGA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RELUGOLIX

---

## Products Affected

- ORGOVYX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# REPOTRECTINIB

---

## Products Affected

- AUGTYRO ORAL CAPSULE 160 MG, 40 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RESMETIROM

## Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	NONALCOHOLIC STEATOHEPATITIS (NASH): INITIAL: DIAGNOSIS CONFIRMED BY BIOPSY OR NONINVASIVE TESTING, OBTAINED IN THE PAST 12 MONTHS, DEMONSTRATING: 1) LIVER FIBROSIS STAGE 2 OR 3, OR 2) NONALCOHOLIC FATTY LIVER DISEASE (NAFLD) ACTIVITY SCORE OF 4 OR MORE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NASH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST, GASTROENTEROLOGIST, OR ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	NASH: RENEWAL: CONTINUES TO HAVE NONCIRRHOTIC NASH WITH MODERATE TO ADVANCED LIVER FIBROSIS (CONSISTENT WITH STAGES F2 TO F3 FIBROSIS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RETIFANLIMAB-DLWR

---

## Products Affected

- ZYNYZ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# REVUMENIB

---

## Products Affected

- REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RIBOCICLIB

---

## Products Affected

- KISQALI 200 MG DAILY DOSE
- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RIBOCICLIB-LETROZOLE

---

## Products Affected

- KISQALI FEMARA CO-PACK ORAL  
TABLET 200 MG/DAY(200 MG X 1)-2.5  
MG, 400 MG/DAY(200 MG X 2)-2.5 MG,  
600 MG/DAY(200 MG X 3)-2.5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RIFAXIMIN

---

## Products Affected

- XIFAXAN ORAL TABLET 200 MG, 550 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	TRAVELERS DIARRHEA, HEPATIC ENCEPHALOPATHY (HE): 12 MOS. IBS-D: 8 WKS.
<b>Other Criteria</b>	HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RILONACEPT

## Products Affected

- ARCALYST

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR S100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES.</p> <p>DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE IL1RN GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR), AND 2) ONE OF THE FOLLOWING: PUSTULAR PSORIASIS-LIKE RASHES, OSTEOMYELITIS, ABSENCE OF BACTERIAL OSTEOMYELITIS, ONYCHOMADESIS. RECURRENT PERICARDITIS (RP): TWO OF THE FOLLOWING: CHEST PAIN CONSISTENT WITH PERICARDITIS, PERICARDIAL FRICTION RUB, ECG SHOWING DIFFUSE ST-SEGMENT ELEVATION OR PR-SEGMENT DEPRESSION, NEW OR WORSENING PERICARDIAL EFFUSION.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CAPS, DIRA: LIFETIME. RP: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	CAPS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR CAPS. DIRA: 1) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR DIRA, AND 2) TRIAL OF THE PREFERRED AGENT: KINERET. RP: 1) HAD AN EPISODE OF ACUTE PERICARDITIS, 2) SYMPTOM-FREE FOR 4 TO 6 WEEKS, AND 3) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR RP.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RIMEGEPANT

## Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# RIOCIQUAT

## Products Affected

- ADEMPAS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) (WHO GROUP 4): WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PAH: NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE (PDE) INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PDE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PDE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RIPRETINIB

---

## Products Affected

- QINLOCK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RISANKIZUMAB-RZAA

## Products Affected

- SKYRIZI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# RITUXIMAB AND HYALURONIDASE HUMAN-SQ

## Products Affected

- RITUXAN HYCELA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	FOLLICULAR LYMPHOMA (FL), DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): 1) HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RITUXIMAB-ABBS

## Products Affected

- TRUXIMA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA, PV: 12 MO. CLL: 6 MO.
<b>Other Criteria</b>	INITIAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: RA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# ROPEGINTERFERON ALFA-2B-NJFT

---

## Products Affected

- BESREMI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RUCAPARIB

---

## Products Affected

- RUBRACA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RUXOLITINIB

---

## Products Affected

- JAKAFI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS
<b>Other Criteria</b>	INITIAL: CHRONIC GRAFT VS HOST DISEASE (CGVHD): NO CONCURRENT USE WITH REZUROCK, NIKTIMVO, OR IMBRUVICA. RENEWAL: MYELOFIBROSIS: CONTINUES TO BENEFIT FROM THE MEDICATION. CGVHD: NO CONCURRENT USE WITH REZUROCK, NIKTIMVO, OR IMBRUVICA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SAPROPTERIN

---

## Products Affected

- *javygtor oral tablet, soluble*
- *sapropterin oral tablet, soluble*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 2 MONTHS, RENEWAL 12 MONTHS.
<b>Other Criteria</b>	HYPERPHENYLALANINEMIA (HPA): INITIAL: NO CONCURRENT USE WITH PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NO CONCURRENT USE WITH PALYNZIQ.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SECUKINUMAB SQ

## Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE  
75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: HS: 12 MONTHS, ALL OTHER INDICATIONS: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. AS, NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID. ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: ALL INDICATIONS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SELEXIPAG

## Products Affected

- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR FOR PAH, 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SELINEXOR

---

## Products Affected

- XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (10 MG X 4), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80 MG/WEEK (80 MG X 1), 80MG TWICE WEEK (160 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# SELPERCATINIB

---

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG
- RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SELUMETINIB

---

## Products Affected

- KOSELUGO ORAL CAPSULE 10 MG, 25 MG
- KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG, 7.5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SEVABERTINIB

---

## Products Affected

- HYRNUO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SILDENAFIL TABLET

## Products Affected

- *sildenafil (pulm.hypertension) oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: AGES 18 YEARS OR OLDER: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. AGES 1 TO 17 YEARS: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PAP GREATER THAN 20 MMHG, 2) PCWP OF 15 MMHG OR LESS, AND 3) PVR OF 3 WOOD UNITS OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL/RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA) OR ANY ORGANIC NITRATES IN ANY FORM AND 2) NO CONCURRENT USE WITH GUANYLATE CYCLASE STIMULATORS.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# SIPONIMOD

---

## Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER(FOR 2MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# SIROLIMUS PROTEIN-BOUND

---

## Products Affected

- FYARRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SODIUM OXYBATE-XYREM

## Products Affected

- *sodium oxybate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CATAPLEXY IN NARCOLEPSY, EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: EDS IN NARCOLEPSY: 1) NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT, AND 2) AGES 18 YEARS OR OLDER: TRIAL, FAILURE OF, OR CONTRAINDICATION TO A FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, OR SUNOSI AND ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT. RENEWAL: CATAPLEXY IN NARCOLEPSY, EDS IN NARCOLEPSY: 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# SOFOSBUVIR/VELPATASVIR

## Products Affected

- EPCLUSA ORAL PELLETS IN PACKET  
150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANA VIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HCV RNA LEVEL WITHIN PAST 6 MONTHS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	No

# SOMATROPIN - NORDITROPIN

## Products Affected

- NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES.
<b>Required Medical Information</b>	INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS BELOW THE MEAN HEIGHT FOR CHILDREN OF THE SAME AGE AND GENDER. TURNER SYNDROME (TS): CONFIRMED BY CHROMOSOMAL ANALYSIS (KARYOTYPING). PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS OF PWS. ADULT GHD: 1) HAS A CONGENITAL, GENETIC, OR ORGANIC DISEASE (E.G., CRANIOPHARYNGIOMA, PITUITARY HYPOPLASIA, ECTOPIC POSTERIOR PITUITARY, PREVIOUS CRANIAL IRRADIATION), OR 2) GHD CONFIRMED BY ONE OF THE FOLLOWING GROWTH HORMONE (GH) STIMULATION TESTS: (A) INSULIN TOLERANCE TEST (PEAK GH OF 5 NG/ML OR LESS), (B) GLUCAGON-STIMULATION TEST (ONE OF THE FOLLOWING: (I) PEAK RESPONSE OF 3 NG/ML OR LESS AND BMI LESS THAN 25 KG/M2, (II) PEAK RESPONSE OF 3 NG/ML OR LESS AND BMI IS BETWEEN 25 - 30 KG/M2 WITH A PRE-TEST PROBABILITY, (III) PEAK RESPONSE OF 1 NG/ML OR LESS AND BMI IS BETWEEN 25 - 30 KG/M2 WITH LOW TEST PROBABILITY, OR (IV) PEAK RESPONSE OF 1 NG/ML OR LESS AND BMI IS GREATER THAN 30 KG/M2), OR (C) MACIMORELIN TEST (PEAK GH OF 2.8 NG/ML OR LESS).
<b>Age Restrictions</b>	SGA: 2 YEARS OF AGE OR OLDER.
<b>Prescriber Restrictions</b>	INITIAL/RENEWAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	<p>INITIAL: PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. INITIAL/RENEWAL: ADULT GHD, PEDIATRIC GHD, SGA, TS, PWS, NOONAN SYNDROME: NO CONCURRENT USE WITH INCRELEX. RENEWAL: ISS: 1) IMPROVEMENT WHILE ON THERAPY (INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PEDIATRIC GHD, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR HAS NOT COMPLETED PREPUBERTAL GROWTH. PWS: IMPROVEMENT IN BODY COMPOSITION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SOMATROPIN - SEROSTIM

## Products Affected

- SEROSTIM SUBCUTANEOUS RECON  
SOLN 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES
<b>Required Medical Information</b>	INITIAL: HIV/WASTING: ONE OF THE FOLLOWING FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS OR 5% WEIGHT LOSS OVER 6 MONTHS, 2) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 3) BCM LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 4) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND BMI LESS THAN 27 KG PER METER SQUARED, OR 5) BMI LESS THAN 20 KG PER METER SQUARED.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 9 MONTHS.
<b>Other Criteria</b>	HIV/WASTING: RENEWAL: 1) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SONIDEGIB

---

## Products Affected

- ODOMZO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC): BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SORAFENIB

---

## Products Affected

- *sorafenib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SOTATERCEPT-CSRK

## Products Affected

- WINREVAIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: 1) ON BACKGROUND PAH THERAPY (FOR AT LEAST 3 MONTHS) WITH AT LEAST TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: A) ORAL ENDOTHELIN RECEPTOR ANTAGONIST, B) ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR FOR PAH, C) ORAL CGMP STIMULATOR, D) IV/SQ PROSTACYCLIN, OR 2) ON ONE AGENT FROM ONE OF THE ABOVE DRUG CLASSES, AND HAS A CONTRAINDICATION OR INTOLERANCE TO ALL OF THE OTHER DRUG CLASSES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# SOTORASIB

---

## Products Affected

- LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# STIRIPENTOL

---

## Products Affected

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SUNITINIB

---

## Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# TADALAFIL - ADCIRCA, ALYQ

## Products Affected

- *alyq*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL/RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA) OR ANY ORGANIC NITRATES IN ANY FORM, AND 2) NO CONCURRENT USE WITH GUANYLATE CYCLASE STIMULATORS.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TADALAFIL-CIALIS

---

## Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA (BPH).
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	BPH: 1) TRIAL OF ONE ALPHA BLOCKER (E.G., DOXAZOSIN, TERAZOSIN, TAMSULOSIN, ALFUZOSIN), AND 2) TRIAL OF ONE 5-ALPHA-REDUCTASE INHIBITOR (E.G., FINASTERIDE, DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TAFAMIDIS

## Products Affected

- VYNDAMAX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CARDIOMYOPATHY ASSOCIATED WITH WILD TYPE OR HEREDITARY TRANSTHYRETIN-MEDIATED AMYLOIDOSIS (ATTR-CM): INITIAL: 1) NEW YORK HEART ASSOCIATION (NYHA) CLASS I, II, OR III HEART FAILURE, AND 2) DIAGNOSIS CONFIRMED BY (A) BONE SCAN (SCINTIGRAPHY) STRONGLY POSITIVE FOR MYOCARDIAL UPTAKE OF TC-99M-PYP, OR (B) BIOPSY OF TISSUE OF AFFECTED ORGAN(S) (CARDIAC AND POSSIBLY NON-CARDIAC SITES) TO CONFIRM AMYLOID PRESENCE AND CHEMICAL TYPING TO CONFIRM PRESENCE OF TRANSTHYRETIN (TTR) PROTEIN.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ATTR-CM: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ATTR SPECIALIST, OR MEDICAL GENETICIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	ATTR-CM: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER ATTR-CM TTR STABILIZERS (E.G., ACORAMIDIS)
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TALAZOPARIB

## Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: 1) HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING, AND 2) IF HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER, RECEIVED PRIOR TREATMENT WITH ENDOCRINE THERAPY OR IS CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# TALETRECTINIB

---

## Products Affected

- IBTROZI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TALQUETAMAB-TGVS

---

## Products Affected

- TALVEY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TARLATAMAB-DLLE

---

## Products Affected

- IMDELLTRA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TAZEMETOSTAT

---

## Products Affected

- TAZVERIK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TEBENTAFUSP-TEBN

---

## Products Affected

- KIMMTRAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TECLISTAMAB-CQYV

---

## Products Affected

- TECVAYLI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TELISOTUZUMAB VEDOTIN-TLLV

---

## Products Affected

- EMRELIS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TELOTRISTAT

---

## Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TEPOTINIB

---

## Products Affected

- TEPMETKO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TERIPARATIDE

---

## Products Affected

- *teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 MONTHS
Other Criteria	OSTEOPOROSIS: HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY, UNLESS REMAINS AT OR HAS RETURNED TO HAVING A HIGH RISK FOR FRACTURE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TESTOSTERONE

## Products Affected

- *testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %)* (25 mg/2.5gram), 1 % (50 mg/5 gram)
- *testosterone transdermal gel in packet 1 %*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TESTOSTERONE CYPIONATE - DEPO

## Products Affected

- *testosterone cypionate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TESTOSTERONE ENANTHATE

## Products Affected

- *testosterone enanthate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: MALE DELAYED PUBERTY: 6MO, MALE HYPOGONADISM: 12 MO. OTHER INDICATIONS: 12 MO.
<b>Other Criteria</b>	INITIAL: MALE HYPOGONADISM: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: MALE HYPOGONADISM: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. MALE DELAYED PUBERTY: HAS NOT RECEIVED MORE THAN TWO 6-MONTH COURSES OF TESTOSTERONE REPLACEMENT THERAPY
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TETRABENAZINE

---

## Products Affected

- *tetrabenazine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# THALIDOMIDE

---

## Products Affected

- THALOMID ORAL CAPSULE 100 MG, 150 MG, 200 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TISLELIZUMAB-JSGR

---

## Products Affected

- TEVIMBRA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TISOTUMAB VEDOTIN-TFTV

---

## Products Affected

- TIVDAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TIVOZANIB

---

## Products Affected

- FOTIVDA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TOCILIZUMAB-AAZG IV

---

## Products Affected

- TYENNE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. GIANT CELL ARTERITIS (GCA): 1) HAS COMPLETED, STARTED, OR WILL SOON START A TAPERING COURSE OF GLUCOCORTICIDS, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: RINVOQ. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ. RENEWAL: RA, PJIA, SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION. GCA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TOCILIZUMAB-AAZG SQ

## Products Affected

- TYENNE
- TYENNE AUTOINJECTOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. GIANT CELL ARTERITIS (GCA): 1) HAS COMPLETED, STARTED, OR WILL SOON START A TAPERING COURSE OF GLUCOCORTICIDS, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: RINVOQ. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TOFACITINIB

## Products Affected

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, PSA, AS, PCJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TOLVAPTAN

## Products Affected

- JYNARQUE ORAL TABLET
- *tolvaptan (polycyst kidney dis) oral tablets, sequential*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	AUTOSOMAL DOMINANT POLYCYSTIC KIDNEY DISEASE (ADPKD): INITIAL: CONFIRMED POLYCYSTIC KIDNEY DISEASE VIA CT, MRI, OR ULTRASOUND.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ADPKD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ADPKD: INITIAL: DOES NOT HAVE ESRD (I.E., RECEIVING DIALYSIS). RENEWAL: HAS NOT PROGRESSED TO ESRD/DIALYSIS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TOPICAL TRETINOIN

---

## Products Affected

- ALTRENO
- *tretinoin topical cream*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TORIPALIMAB-TPZI

---

## Products Affected

- LOQTORZI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	NASOPHARYNGEAL CARCINOMA (NPC): FIRST LINE TREATMENT: 24 MOS, PREVIOUSLY TREATED: LIFETIME.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TOVORAFENIB

---

## Products Affected

- OJEMDA ORAL SUSPENSION FOR RECONSTITUTION
- OJEMDA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TRAMETINIB SOLUTION

## Products Affected

- MEKINIST ORAL RECON SOLN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE OR METASTATIC MELANOMA, MELANOMA, METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), LOCALLY ADVANCED OR METASTATIC ANAPLASTIC THYROID CANCER (ATC), UNRESECTABLE OR METASTATIC SOLID TUMOR, LOW-GRADE GLIOMA (LGG): UNABLE TO SWALLOW MEKINIST TABLETS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TRAMETINIB TABLET

---

## Products Affected

- MEKINIST ORAL TABLET 0.5 MG, 2 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TRASTUZUMAB-DKST

---

## Products Affected

- OGIVRI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TRASTUZUMAB-HYALURONIDASE-OYSK

---

## Products Affected

- HERCEPTIN HYLECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# TRAZODONE

---

## Products Affected

- RALDESY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MAJOR DEPRESSIVE DISORDER (MDD): CONTRAINDICATION TO OR UNABLE TO SWALLOW TRAZODONE TABLETS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TREMELIMUMAB-ACTL

---

## Products Affected

- IMJUDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS.
Other Criteria	UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO. NSCLC: HAS NOT RECEIVED A TOTAL OF 5 DOSES OF IMJUDO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TRIENTINE CAPSULE

## Products Affected

- *trientine oral capsule 250 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	WILSONS DISEASE: INITIAL: LEIPZIG SCORE OF 4 OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
<b>Other Criteria</b>	WILSONS DISEASE: INITIAL: TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE TABLET. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# TRIFLURIDINE/TIPIRACIL

---

## Products Affected

- LONSURF ORAL TABLET 15-6.14 MG,  
20-8.19 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TRIPTORELIN-TRELSTAR

---

## Products Affected

- TRELSTAR INTRAMUSCULAR  
SUSPENSION FOR RECONSTITUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TUCATINIB

---

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# UBROGEPANT

## Products Affected

- UBRELVY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ACUTE MIGRAINE TREATMENT: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN), AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT. RENEWAL: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT, AND 2) ONE OF THE FOLLOWING: (A) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR (B) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# UPADACITINIB

---

## Products Affected

- RINVOQ
- RINVOQ LQ

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. ULCERATIVE COLITIS (UC), CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE CONVENTIONAL SYNTHETIC DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE OF AT LEAST 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. AD: 1) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, AND 2) TRIAL OF OR CONTRAINDICATION TO A TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR. AS, NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). GIANT CELL ARTERITIS (GCA): HAS COMPLETED, STARTED, OR WILL SOON START A TAPERING COURSE OF GLUCOCORTICIDS.</p> <p>INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: RA, PSA, AS, NR-AXSPA, PJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. AD: IMPROVEMENT WHILE ON THERAPY.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# USTEKINUMAB-AAUZ SQ

## Products Affected

- *ustekinumab-aauz*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# USTEKINUMAB-AEKN IV

## Products Affected

- SELARSDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# USTEKINUMAB-AEKN SQ

## Products Affected

- SELARSDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# USTEKINUMAB-KFCE IV

## Products Affected

- YESINTEK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# USTEKINUMAB-KFCE SQ

## Products Affected

- YESINTEK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSO: 1) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY), 2) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA, OR 3) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION. INITIAL/RENEWAL: ALL INDICATIONS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR THE SAME INDICATION. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VALBENAZINE

---

## Products Affected

- INGREZZA
- INGREZZA INITIATION PK(TARDIV)
- INGREZZA SPRINKLE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	TARDIVE DYSKINESIA (TD): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST. CHOREA ASSOCIATED WITH HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TD: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VANDETANIB

---

## Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VANZACAFTOR-TEZACAFTOR- DEUTIVACAFTOR

## Products Affected

- ALYFTREK ORAL TABLET 10-50-125  
MG, 4-20-50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
Other Criteria	CF: INITIAL: NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR. RENEWAL: 1) IMPROVEMENT IN CLINICAL STATUS, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VEMURAFENIB

---

## Products Affected

- ZELBORAF

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VENETOCLAX

---

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VERICIGUAT

## Products Affected

- VERQUVO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL:12 MONTHS.
<b>Other Criteria</b>	HEART FAILURE (HF): INITIAL: 1) TRIAL OF, CONTRAINDICATION, OR INTOLERANCE TO ONE PREFERRED SGLT-2 INHIBITOR, AND 2) TRIAL OF, CONTRAINDICATION, OR INTOLERANCE TO ONE AGENT FROM ANY OF THE FOLLOWING STANDARD OF CARE CLASSES: (A) ACE INHIBITOR, ARB, OR ARNI, (B) BETA BLOCKER (BISOPROLOL, CARVEDILOL, METOPROLOL SUCCINATE), OR (C) ALDOSTERONE ANTAGONIST (SPIRONOLACTONE, EPLERENONE). INITIAL/RENEWAL: NO CONCURRENT USE WITH RIOCIGUAT OR PDE-5 INHIBITORS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VIGABATRIN

---

## Products Affected

- *vigabatin*
- *vigadrone*
- *vigpoder*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: CPS: TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# VIMSELTINIB

---

## Products Affected

- ROMVIMZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VISMODEGIB

---

## Products Affected

- ERIVEDGE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VONOPRAZAN

## Products Affected

- VOQUEZNA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: EROSIIVE ESOPHAGITIS (EE): DIAGNOSIS CONFIRMED BY ENDOSCOPY (E.G., LOS ANGELES CLASSIFICATION OF REFLUX ESOPHAGITIS GRADE A-D). NON-EROSIVE GASTROESOPHAGEAL REFLUX DISEASE (NERD): DIAGNOSIS CONFIRMED BY ENDOSCOPY AND DOES NOT HAVE PRESENCE OF VISIBLE EROSION (E.G., DOES NOT HAVE LOS ANGELES CLASSIFICATION OF REFLUX ESOPHAGITIS GRADE A-D).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: EE, NERD: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: H PYLORI: 30 DAYS. EE: 8 WEEKS. NERD: 4 WEEKS. RENEWAL: EE: 24 WEEKS.
<b>Other Criteria</b>	INITIAL: EE: TRIAL OF OR CONTRAINDICATION TO TWO PROTON PUMP INHIBITORS AT MAXIMUM DOSE FOR 8 WEEKS EACH. NERD: 1) NO PREVIOUS TREATMENT FAILURE WITH VOQUEZNA IN THE LAST 12 MONTHS, AND 2) TRIAL OF OR CONTRAINDICATION TO TWO PROTON PUMP INHIBITORS AT MAXIMUM DOSE FOR 8 WEEKS EACH. RENEWAL: EE: MAINTAINED A CLINICAL RESPONSE ON VOQUEZNA
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prerequisite Therapy Required</b>	Yes

# VORASIDENIB

---

## Products Affected

- VORANIGO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VORICONAZOLE SUSPENSION

---

## Products Affected

- *voriconazole oral suspension for reconstitution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CANDIDA INFECTIONS: 3 MOS. CONTINUATION OF THERAPY, ALL OTHER INDICATIONS: 6 MOS.
<b>Other Criteria</b>	CANDIDA INFECTIONS: CONTRAINDICATION TO OR UNABLE TO SWALLOW FLUCONAZOLE TABLETS. ALL INDICATIONS EXCEPT ESOPHAGEAL CANDIDIASIS: UNABLE TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ZANIDATAMAB-HR11

---

## Products Affected

- ZIIHERA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ZANUBRUTINIB

---

## Products Affected

- BRUKINSA ORAL CAPSULE
- BRUKINSA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MANTLE CELL LYMPHOMA: INTOLERANCE TO CALQUENCE. CHRONIC LYMPHOCYTIC LEUKEMIA, SMALL LYMPHOCYTIC LYMPHOMA: INTOLERANCE TO CALQUENCE OR IMBRUVICA. WALDENSTROMS MACROGLOBULINEMIA: NO STEP REQUIRED.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ZENOCUTUZUMAB-ZBCO

---

## Products Affected

- BIZENGRI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ZOLBETUXIMAB-CLZB

---

## Products Affected

- VYLOY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ZONGERTINIB

---

## Products Affected

- HERNEXEOS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ZURANOLONE

---

## Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	14 DAYS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

## INDEX

1ST TIER UNIFINE PENTP 5MM 31G... 166	AQINJECT PEN NEEDLE 32G 4MM..... 166
1ST TIER UNIFINE PNTIP 4MM 32G.....166	ARCALYST.....287
1ST TIER UNIFINE PNTIP 6MM 31G.....166	ARIKAYCE.....21
1ST TIER UNIFINE PNTIP 8MM 31G	<i>armodafinil</i> .....228
STRL,SINGLE-USE,SHRT..... 166	ASSURE ID DUO PRO NDL 31G 5MM.. 166
1ST TIER UNIFINE PNTIP 29GX1/2"..... 166	ASSURE ID DUO-SHIELD 30GX3/16" ... 166
1ST TIER UNIFINE PNTIP 31GX3/16..... 166	ASSURE ID DUO-SHIELD 30GX5/16" ... 166
1ST TIER UNIFINE PNTIP 32GX5/32..... 166	ASSURE ID INSULIN SAFETY
<i>abiraterone</i> ..... 7	SYRINGE 1 ML 29 GAUGE X 1/2"..... 166
<i>abiraterone, submicronized</i> ..... 8	ASSURE ID PEN NEEDLE 30GX3/16" ... 166
<i>abirtega</i> .....7	ASSURE ID PEN NEEDLE 30GX5/16" ... 166
ABOUTTIME PEN NEEDLE.....166	ASSURE ID PEN NEEDLE 31GX3/16" ... 166
ACTIMMUNE.....179	ASSURE ID PRO PEN NDL 30G 5MM... 166
ADEMPAS.....291	ASSURE ID SYR 0.5 ML 31GX15/64" .... 166
ADVOCATE INS 0.3 ML 30GX5/16" ..... 166	ASSURE ID SYR 1 ML 31GX15/64" ..... 166
ADVOCATE INS 0.3 ML 31GX5/16" ..... 166	ATTRUBY..... 10
ADVOCATE INS 0.5 ML 30GX5/16"..... 166	AUGTYRO ORAL CAPSULE 160 MG,
ADVOCATE INS 0.5 ML 31GX5/16" ..... 166	40 MG.....280
ADVOCATE INS 1 ML 31GX5/16"..... 166	AUSTEDO ORAL TABLET 12 MG, 6
ADVOCATE INS SYR 0.3 ML 29GX1/2. 166	MG, 9 MG..... 89
ADVOCATE INS SYR 0.5 ML 29GX1/2. 166	AUSTEDO XR ORAL TABLET
ADVOCATE INS SYR 1 ML 29GX1/2" .. 166	EXTENDED RELEASE 24 HR 12 MG, 18
ADVOCATE INS SYR 1 ML 30GX5/16.. 166	MG, 24 MG, 30 MG, 36 MG, 42 MG, 48
ADVOCATE PEN NDL 12.7MM 29G.... 166	MG, 6 MG..... 89
ADVOCATE PEN NEEDLE 32G 4MM... 166	AUSTEDO XR TITRATION KT(WK1-4)..89
ADVOCATE PEN NEEDLE 4MM 33G... 166	AUTOSHIELD DUO PEN NDL 30G
ADVOCATE PEN NEEDLES 5MM 31G. 166	5MM..... 166
ADVOCATE PEN NEEDLES 8MM 31G. 166	AVMAPKI.....39
AIMOVIG AUTOINJECTOR..... 122	AVMAPKI-FAKZYNJA.....39
AKEEGA.....236	AVONEX INTRAMUSCULAR PEN
ALCOHOL PADS.....166	INJECTOR KIT.....176
ALCOHOL PREP SWABS.....166	AVONEX INTRAMUSCULAR
ALCOHOL WIPES..... 166	SYRINGE KIT..... 176
ALECENSA..... 19	AYVAKIT.....38
ALTRENO.....360	BALVERSA ORAL TABLET 3 MG, 4
ALUNBRIG ORAL TABLET 180 MG, 30	MG, 5 MG..... 121
MG, 90 MG..... 58	BD AUTOSHIELD DUO NDL
ALUNBRIG ORAL TABLETS,DOSE	5MMX30G..... 166
PACK..... 58	BD ECLIPSE 30GX1/2" SYRINGE..... 166
ALVAIZ..... 109	BD ECLIPSE NEEDLE 30GX1/2" (OTC) 166
ALYFTREK ORAL TABLET 10-50-125	BD INS SYR 0.3 ML 8MMX31G(1/2).....166
MG, 4-20-50 MG.....388	BD INS SYR UF 0.3 ML 12.7MMX30G.. 166
<i>alyq</i> ..... 330	BD INS SYR UF 0.5 ML 12.7MMX30G
ANKTIVA.....242	NOT FOR RETAIL SALE..... 166
AQINJECT PEN NEEDLE 31G 5MM..... 166	BD INSULIN SYR 1 ML 27GX12.7MM.. 166

BD INSULIN SYR 1 ML 27GX5/8" MICRO-FINE.....	166	CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG.....	61
BD LO-DOSE ULTRA-FINE.....	166	CALQUENCE.....	9
BD NANO 2 GEN PEN NDL 32G 4MM..	166	CALQUENCE (ACALABRUTINIB MAL).....	9
BD SAFETGLD INS 0.3 ML 29G 13MM.	166	CAMCEVI (6 MONTH).....	196
BD SAFETYGLD INS 0.3 ML 31G 8MM	166	CAMZYOS.....	214
BD SAFETYGLD INS 0.5 ML 30G 8MM	166	CAPRELSA ORAL TABLET 100 MG, 300 MG.....	387
BD SAFETYGLD INS 1 ML 29G 13MM.	166	CAREFINE PEN NEEDLE 12.7MM 29G.	166
BD SAFETYGLID INS 1 ML 6MMX31G	166	CAREFINE PEN NEEDLE 4MM 32G.....	166
BD SAFETYGLIDE SYRINGE 27GX5/8	166	CAREFINE PEN NEEDLE 5MM 32G.....	166
BD SAFTYGLD INS 0.3 ML 6MMX31G	166	CAREFINE PEN NEEDLE 6MM 31G.....	166
BD SAFTYGLD INS 0.5 ML 29G 13MM	166	CAREFINE PEN NEEDLE 8MM 30G.....	166
BD SAFTYGLD INS 0.5 ML 6MMX31G	166	CAREFINE PEN NEEDLES 6MM 32G...	166
BD SINGLE USE SWAB.....	166	CAREFINE PEN NEEDLES 8MM 31G...	166
BD UF MICRO PEN NEEDLE 6MMX32G.....	166	CARETOUCH ALCOHOL 70% PREP PAD.....	166
BD UF MINI PEN NEEDLE 5MMX31G.	166	CARETOUCH PEN NEEDLE 29G 12MM .....	166
BD UF NANO PEN NEEDLE 4MMX32G .....	166	CARETOUCH PEN NEEDLE 31GX1/4".	166
BD UF ORIG PEN NDL 12.7MMX29G...	166	CARETOUCH PEN NEEDLE 31GX3/16" .....	166
BD UF SHORT PEN NEEDLE 8MMX31G.....	166	CARETOUCH PEN NEEDLE 31GX5/16" .....	166
BD VEO INS 0.3 ML 6MMX31G (1/2)....	166	CARETOUCH PEN NEEDLE 32GX3/16" .....	166
BD VEO INS SYRING 1 ML 6MMX31G	166	CARETOUCH PEN NEEDLE 32GX5/32" .....	166
BD VEO INS SYRN 0.3 ML 6MMX31G.	166	CARETOUCH SYR 0.3 ML 31GX5/16" ..	166
BD VEO INS SYRN 0.5 ML 6MMX31G.	166	CARETOUCH SYR 0.5 ML 30GX5/16" ..	166
<i>bendamustine intravenous recon soln</i> .....	48	CARETOUCH SYR 0.5 ML 31GX5/16" ..	166
BENDAMUSTINE INTRAVENOUS SOLUTION.....	48	CARETOUCH SYR 1 ML 28GX5/16" ..	166
BENDEKA.....	48	CARETOUCH SYR 1 ML 29GX5/16" ..	166
BENLYSTA SUBCUTANEOUS.....	45	CARETOUCH SYR 1 ML 30GX5/16" ..	166
BESREMI.....	299	CARETOUCH SYR 1 ML 31GX5/16" ..	166
<i>betaine</i> .....	51	<i>carglumic acid</i> .....	65
BETASERON SUBCUTANEOUS KIT ....	177	CAYSTON.....	43
<i>bexarotene</i> .....	53	CIMZIA 200 MG/ML SYRINGE KIT.....	67
BIZENGRI.....	401	CIMZIA POWDER FOR RECONST.....	67
BORDERED GAUZE 2"X2".....	166	CIMZIA STARTER KIT.....	67
<i>bortezomib injection</i> .....	55	CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2).....	67
BORUZU.....	55	CLICKFINE PEN NEEDLE 32GX5/32" 32GX4MM, STERILE.....	166
<i>bosentan oral tablet</i> .....	56		
BOSULIF ORAL CAPSULE 100 MG, 50 MG.....	57		
BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG.....	57		
BRAFTOVI.....	113		
BRUKINSA ORAL CAPSULE.....	400		
BRUKINSA ORAL TABLET.....	400		

COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY).....	60	COMFORT EZ SYR 1 ML 28GX1/2" .....	166
COMFORT EZ 0.3 ML 31G 15/64" .....	166	COMFORT EZ SYR 1 ML 29GX1/2" .....	166
COMFORT EZ 0.5 ML 31G 15/64" .....	166	COMFORT EZ SYR 1 ML 30GX1/2" .....	166
COMFORT EZ INS 0.3 ML 30GX1/2" .....	166	COMFORT EZ SYR 1 ML 30GX5/16" .....	166
COMFORT EZ INS 0.3 ML 30GX5/16" .....	166	COMFORT POINT PEN NDL 31GX1/3" .....	166
COMFORT EZ INS 1 ML 31G 15/64" .....	166	COMFORT POINT PEN NDL 31GX1/6" .....	166
COMFORT EZ INS 1 ML 31GX5/16" .....	166	COMFORT TOUCH PEN NDL 31G 4MM .....	166
COMFORT EZ INSULIN SYR 0.3 ML....	166	COMFORT TOUCH PEN NDL 31G 5MM .....	166
COMFORT EZ INSULIN SYR 0.5 ML....	166	COMFORT TOUCH PEN NDL 31G 6MM .....	166
COMFORT EZ PEN NEEDLE 12MM 29G.....	166	COMFORT TOUCH PEN NDL 31G 8MM .....	166
COMFORT EZ PEN NEEDLES 4MM 32G SINGLE USE, MICRO.....	166	COMFORT TOUCH PEN NDL 32G 4MM .....	166
COMFORT EZ PEN NEEDLES 4MM 33G.....	166	COMFORT TOUCH PEN NDL 32G 5MM .....	166
COMFORT EZ PEN NEEDLES 5MM 31G MINI.....	166	COMFORT TOUCH PEN NDL 32G 6MM .....	166
COMFORT EZ PEN NEEDLES 5MM 32G SINGLE USE,MINI,HRI.....	166	COMFORT TOUCH PEN NDL 32G 8MM .....	166
COMFORT EZ PEN NEEDLES 5MM 33G.....	166	COMFORT TOUCH PEN NDL 33G 4MM .....	166
COMFORT EZ PEN NEEDLES 6MM 31G.....	166	COMFORT TOUCH PEN NDL 33G 6MM .....	166
COMFORT EZ PEN NEEDLES 6MM 32G.....	166	COMFORT TOUCH PEN NDL 33GX5MM.....	166
COMFORT EZ PEN NEEDLES 6MM 33G.....	166	COPIKTRA .....	101
COMFORT EZ PEN NEEDLES 8MM 31G SHORT.....	166	CORTROPHIN GEL INJECTION.....	73
COMFORT EZ PEN NEEDLES 8MM 32G.....	166	COSENTYX (2 SYRINGES).....	303
COMFORT EZ PEN NEEDLES 8MM 33G.....	166	COSENTYX PEN (2 PENS).....	303
COMFORT EZ PRO PEN NDL 30G 8MM .....	166	COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML .....	303
COMFORT EZ PRO PEN NDL 31G 4MM .....	166	COSENTYX UNOREADY PEN.....	303
COMFORT EZ PRO PEN NDL 31G 5MM .....	166	COTELLIC .....	72
COMFORT EZ SYR 0.3 ML 29GX1/2" .....	166	CRESEMBA ORAL.....	181
COMFORT EZ SYR 0.5 ML 28GX1/2" .....	166	CURAD GAUZE PADS 2" X 2".....	166
COMFORT EZ SYR 0.5 ML 29GX1/2" .....	166	CURITY ALCOHOL PREPS 2 PLY,MEDIUM.....	166
COMFORT EZ SYR 0.5 ML 30GX1/2" .....	166	CURITY GAUZE PADS.....	166
COMFORT EZ SYR 1 ML 27G 12.7MM. ....	166	CURITY GAUZE SPONGES (12 PLY)- 200/BAG.....	166
		CYLTEZO(CF).....	16
		CYLTEZO(CF) PEN.....	16
		CYLTEZO(CF) PEN CROHN'S-UC-HS.....	16
		CYLTEZO(CF) PEN PSORIASIS-UV .....	16

<i>dalfampridine</i> .....	80	DROPLET INS SYR 0.5 ML 30G 8MM...	166
DANYELZA.....	229	DROPLET INS SYR 0.5 ML 31G 6MM...	166
DANZITEN.....	232	DROPLET INS SYR 0.5 ML 31G 8MM...	166
<i>dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg</i> .....	82	DROPLET INS SYR 1 ML 29G 12.7MM.	166
DATROWAY.....	83	DROPLET INS SYR 1 ML 30G 8MM.....	166
DAURISMO ORAL TABLET 100 MG, 25 MG.....	147	DROPLET INS SYR 1 ML 30GX12.5MM.....	166
<i>deferasirox oral granules in packet</i> .....	85	DROPLET INS SYR 1 ML 30GX6MM....	166
<i>deferasirox oral tablet</i> .....	85	DROPLET INS SYR 1 ML 31G 6MM.....	166
DERMACEA 2"X2" GAUZE 12 PLY, USP TYPE VII.....	166	DROPLET INS SYR 1 ML 31GX8MM....	166
DERMACEA GAUZE 2"X2" SPONGE 8 PLY.....	166	DROPLET MICRON 34G X 9/64".....	166
DERMACEA NON-WOVEN 2"X2" SPNGE.....	166	DROPLET PEN NEEDLE 29G 10MM.....	166
<i>dermacinrx lidocan 5% patch outer</i> .....	204	DROPLET PEN NEEDLE 29G 12MM.....	166
DIACOMIT ORAL CAPSULE 250 MG, 500 MG.....	328	DROPLET PEN NEEDLE 30G 8MM.....	166
DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG.....	328	DROPLET PEN NEEDLE 31G 5MM.....	166
<i>diclofenac epolamine</i> .....	91	DROPLET PEN NEEDLE 31G 6MM.....	166
<i>diclofenac sodium topical solution in metered-dose pump</i> .....	90	DROPLET PEN NEEDLE 31G 8MM.....	166
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg</i> .....	92	DROPLET PEN NEEDLE 32G 4MM.....	166
<i>dronabinol</i> .....	96	DROPLET PEN NEEDLE 32G 5MM.....	166
DROPLET 0.3 ML 29G 12.7MM(1/2).....	166	DROPLET PEN NEEDLE 32G 6MM.....	166
DROPLET 0.3 ML 30G 12.7MM(1/2).....	166	DROPLET PEN NEEDLE 32G 8MM.....	166
DROPLET 0.5 ML 29GX12.5MM(1/2).....	166	DROPSAFE ALCOHOL 70% PREP PADS.....	166
DROPLET 0.5 ML 30GX12.5MM(1/2).....	166	DROPSAFE INS SYR 0.3 ML 31G 6MM	166
DROPLET INS 0.3 ML 29GX12.5MM.....	166	DROPSAFE INS SYR 0.3 ML 31G 8MM	166
DROPLET INS 0.3 ML 30G 8MM(1/2)....	166	DROPSAFE INS SYR 0.5 ML 31G 6MM	166
DROPLET INS 0.3 ML 30GX12.5MM.....	166	DROPSAFE INS SYR 0.5 ML 31G 8MM	166
DROPLET INS 0.3 ML 31G 6MM(1/2)....	166	DROPSAFE INSUL SYR 1 ML 31G 6MM.....	166
DROPLET INS 0.3 ML 31G 8MM(1/2)....	166	DROPSAFE INSUL SYR 1 ML 31G 8MM.....	166
DROPLET INS 0.5 ML 29G 12.7MM.....	166	DROPSAFE INSULN 1 ML 29G 12.5MM.....	166
DROPLET INS 0.5 ML 30G 12.7MM.....	166	DROPSAFE INSULN 1 ML 29G 12.5MM.....	166
DROPLET INS 0.5 ML 30GX6MM(1/2)..	166	DROPSAFE PEN NEEDLE 31G 5MM.....	166
DROPLET INS 0.5 ML 30GX8MM(1/2)..	166	DROPSAFE PEN NEEDLE 31G 8MM.....	166
DROPLET INS 0.5 ML 31GX6MM(1/2)..	166	DROPSAFE PEN NEEDLE 31GX1/4".....	166
DROPLET INS 0.5 ML 31GX8MM(1/2)..	166	<i>droxidopa</i> .....	97
DROPLET INS SYR 0.3 ML 30GX6MM.	166	DRUG MART ULTRA COMFORT SYR.	166
DROPLET INS SYR 0.3 ML 30GX8MM.	166	DUPIXENT PEN.....	98
DROPLET INS SYR 0.3 ML 31GX6MM.	166	DUPIXENT SYRINGE.....	98
DROPLET INS SYR 0.3 ML 31GX8MM.	166	EASY CMFT SFTY PEN NDL 31G 5MM	166
		EASY CMFT SFTY PEN NDL 31G 6MM	166
		EASY CMFT SFTY PEN NDL 32G 4MM	166
		EASY COMFORT 0.3 ML 31G 1/2".....	166
		EASY COMFORT 0.3 ML 31G 5/16".....	166
		EASY COMFORT 0.3 ML SYRINGE.....	166
		EASY COMFORT 0.5 ML 30GX1/2".....	166

EASY COMFORT 0.5 ML 31GX5/16" .....	166	EASY TOUCH PEN NEEDLE 29GX1/2"	166
EASY COMFORT 0.5 ML 32GX5/16" .....	166	EASY TOUCH PEN NEEDLE 30GX5/16	166
EASY COMFORT 0.5 ML SYRINGE.....	166	EASY TOUCH PEN NEEDLE 31GX1/4"	166
EASY COMFORT 1 ML 31GX5/16" .....	166	EASY TOUCH PEN NEEDLE 31GX3/16	166
EASY COMFORT 1 ML 32GX5/16" .....	166	EASY TOUCH PEN NEEDLE 31GX5/16	166
EASY COMFORT ALCOHOL 70% PAD	166	EASY TOUCH PEN NEEDLE 32GX1/4"	166
EASY COMFORT INSULIN 1 ML SYR..	166	EASY TOUCH PEN NEEDLE 32GX3/16	166
EASY COMFORT PEN NDL 29G 4MM..	166	EASY TOUCH PEN NEEDLE 32GX5/32	166
EASY COMFORT PEN NDL 29G 5MM..	166	EASY TOUCH SAF PEN NDL 29G 5MM	
EASY COMFORT PEN NDL 31GX1/4" ..	166	.....	166
EASY COMFORT PEN NDL 31GX3/16"	166	EASY TOUCH SAF PEN NDL 29G 8MM	
EASY COMFORT PEN NDL 31GX5/16"	166	.....	166
EASY COMFORT PEN NDL 32GX5/32"	166	EASY TOUCH SAF PEN NDL 30G 5MM	
EASY COMFORT PEN NDL 33G 4MM..	166	.....	166
EASY COMFORT PEN NDL 33G 5MM..	166	EASY TOUCH SAF PEN NDL 30G 8MM	
EASY COMFORT PEN NDL 33G 6MM..	166	.....	166
EASY COMFORT SYR 0.5 ML 29G		EASY TOUCH SYR 0.5 ML 28G	
8MM.....	166	12.7MM.....	166
EASY COMFORT SYR 1 ML 29G 8MM.	166	EASY TOUCH SYR 0.5 ML 29G	
EASY COMFORT SYR 1 ML 30GX1/2" .	166	12.7MM.....	166
EASY GLIDE INS 0.3 ML 31GX6MM....	166	EASY TOUCH SYR 1 ML 27G 16MM....	166
EASY GLIDE INS 0.5 ML 31GX6MM....	166	EASY TOUCH SYR 1 ML 28G 12.7MM.	166
EASY GLIDE INS 1 ML 31GX6MM.....	166	EASY TOUCH SYR 1 ML 29G 12.7MM.	166
EASY GLIDE PEN NEEDLE 4MM 33G..	166	EASY TOUCH UNI-SLIP SYR 1 ML.....	166
EASY TOUCH 0.3 ML SYR 30GX1/2" ....	166	EASYTOUCH SAF PEN NDL 30G 6MM	166
EASY TOUCH 0.5 ML SYR 27GX1/2" ....	166	ELAHERE.....	225
EASY TOUCH 0.5 ML SYR 29GX1/2" ....	166	ELIGARD.....	197
EASY TOUCH 0.5 ML SYR 30GX1/2" ....	166	ELIGARD (3 MONTH).....	197
EASY TOUCH 0.5 ML SYR 30GX5/16...	166	ELIGARD (4 MONTH).....	197
EASY TOUCH 1 ML SYR 27GX1/2" .....	166	ELIGARD (6 MONTH).....	197
EASY TOUCH 1 ML SYR 29GX1/2" .....	166	ELREXFIO 44 MG/1.1 ML VIAL INNER,	
EASY TOUCH 1 ML SYR 30GX1/2" .....	166	SUV, P/F.....	108
EASY TOUCH ALCOHOL 70% PADS		ELREXFIO SUBCUTANEOUS	
GAMMA-STERILIZED.....	166	SOLUTION 40 MG/ML.....	108
EASY TOUCH FLIPLOK 1 ML 27GX0.5	166	<i>eltrombopag olamine oral powder in</i>	
EASY TOUCH INSULIN 1 ML 29GX1/2	166	<i>packet 12.5 mg, 25 mg.....</i>	110
EASY TOUCH INSULIN 1 ML 30GX1/2	166	<i>eltrombopag olamine oral tablet 12.5 mg,</i>	
EASY TOUCH INSULIN SYR 0.3 ML....	166	<i>25 mg, 50 mg, 75 mg.....</i>	110
EASY TOUCH INSULIN SYR 0.5 ML....	166	EMBRACE PEN NEEDLE 29G 12MM....	166
EASY TOUCH INSULIN SYR 1 ML.....	166	EMBRACE PEN NEEDLE 30G 5MM.....	166
EASY TOUCH INSULIN SYR 1 ML		EMBRACE PEN NEEDLE 30G 8MM.....	166
RETRACTABLE.....	166	EMBRACE PEN NEEDLE 31G 5MM.....	166
EASY TOUCH INSULN 1 ML 29GX1/2"	166	EMBRACE PEN NEEDLE 31G 6MM.....	166
EASY TOUCH INSULN 1 ML 30GX1/2"	166	EMBRACE PEN NEEDLE 31G 8MM.....	166
EASY TOUCH INSULN 1 ML 30GX5/16	166	EMBRACE PEN NEEDLE 32G 4MM.....	166
EASY TOUCH INSULN 1 ML 31GX5/16	166	EMGALITY PEN.....	142
EASY TOUCH LUER LOK INSUL 1 ML	166		

EMGALITY SYRINGE	
SUBCUTANEOUS SYRINGE 120	
MG/ML, 300 MG/3 ML (100 MG/ML X	
3).....	142
EMRELIS.....	341
ENBREL.....	125
ENBREL MINI.....	125
ENBREL SURECLICK.....	125
ENSACOVE ORAL CAPSULE 100 MG,	
25 MG.....	114
EPCLUSA ORAL PELLETS IN PACKET	
150-37.5 MG, 200-50 MG.....	316
EPCLUSA ORAL TABLET.....	316
EPIDIOLEX.....	62
EPKINLY.....	118
EQL INSULIN 0.5 ML SYRINGE.....	166
EQL INSULIN 0.5 ML SYRINGE	
SHORT NEEDLE.....	166
ERBITUX.....	69
ERIVEDGE.....	394
ERLEADA ORAL TABLET 240 MG, 60	
MG.....	26
<i>erlotinib oral tablet 100 mg, 150 mg, 25</i>	
<i>mg.....</i>	123
<i>everolimus (antineoplastic) oral tablet 10</i>	
<i>mg, 2.5 mg, 5 mg, 7.5 mg.....</i>	127
<i>everolimus (antineoplastic) oral tablet for</i>	
<i>suspension.....</i>	128
EXEL U100 0.3 ML 29GX1/2".....	166
EXXUA ORAL TABLET EXTENDED	
RELEASE 24 HR.....	145
FAKZYNJA.....	39
FASENRA.....	49
FASENRA PEN.....	49
<i>fentanyl citrate buccal lozenge on a handle</i>	
<i>.....</i>	132
<i>fingolimod.....</i>	138
FINTEPLA.....	131
FOTIVDA.....	352
FP INSULIN 1 ML SYRINGE.....	166
FREESTYLE PREC 0.5 ML 30GX5/16....	166
FREESTYLE PREC 0.5 ML 31GX5/16....	166
FREESTYLE PREC 1 ML 30GX5/16".....	166
FREESTYLE PREC 1 ML 31GX5/16".....	166
FRUZAQLA ORAL CAPSULE 1 MG, 5	
MG.....	140
FYARRO.....	313
GAUZE PAD TOPICAL BANDAGE 2 X	
2 ".....	166
GAUZE PADS 2"X2" STRL.....	166
GAVRETO.....	274
<i>gefitinib.....</i>	144
GILOTRIF.....	18
<i>glatiramer subcutaneous syringe 20 mg/ml,</i>	
<i>40 mg/ml.....</i>	148
<i>glatopa subcutaneous syringe 20 mg/ml, 40</i>	
<i>mg/ml.....</i>	148
<i>glutamine (sickle cell).....</i>	202
GNP ALCOHOL SWAB STERILE, TWO	
PLY.....	166
GNP CLICKFINE 31G X 1/4" NDL 6MM,	
UNIVERSAL.....	166
GNP CLICKFINE 31G X 5/16" NDL	
8MM, UNIVERSAL.....	166
GNP PEN NEEDLE 31G 5MM.....	166
GNP PEN NEEDLE 32G 4MM.....	166
GNP PEN NEEDLE 32G 6MM.....	166
GNP SIMPLI PEN NEEDLE 32G 4MM...	166
GNP ULT C 0.3 ML 29GX1/2" (1/2) 1/2	
UNIT.....	166
GNP ULT CMFRT 0.5 ML 29GX1/2".....	166
GNP ULTRA COMFORT 0.5 ML SYR....	166
GNP ULTRA COMFORT 1 ML	
SYRINGE.....	166
GNP ULTRA COMFORT 3/10 ML SYR..	166
GOMEKLI ORAL CAPSULE 1 MG, 2	
MG.....	224
GOMEKLI ORAL TABLET FOR	
SUSPENSION.....	224
GS PEN NEEDLE 31G X 8MM.....	166
HAEGARDA SUBCUTANEOUS RECON	
SOLN 2,000 UNIT, 3,000 UNIT.....	59
HARVONI ORAL PELLETS IN PACKET	
33.75-150 MG, 45-200 MG.....	190
HARVONI ORAL TABLET.....	190
HEALTHWISE INS 0.3 ML 30GX5/16"..	166
HEALTHWISE INS 0.3 ML 31GX5/16"..	166
HEALTHWISE INS 0.5 ML 30GX5/16"..	166
HEALTHWISE INS 0.5 ML 31GX5/16"..	166
HEALTHWISE INS 1 ML 30GX5/16".....	166
HEALTHWISE INS 1 ML 31GX5/16".....	166
HEALTHWISE PEN NEEDLE 31G 5MM	166
HEALTHWISE PEN NEEDLE 31G 8MM	166
HEALTHWISE PEN NEEDLE 32G 4MM	166

HEALTHY ACCENTS PENTIP 4MM 32G.....	166	INGREZZA.....	386
HEALTHY ACCENTS PENTIP 5MM 31G.....	166	INGREZZA INITIATION PK(TARDIV)..	386
HEALTHY ACCENTS PENTIP 6MM 31G.....	166	INGREZZA SPRINKLE.....	386
HEALTHY ACCENTS PENTIP 8MM 31G.....	166	INLURIYO.....	162
HEALTHY ACCENTS PENTIP 12MM 29G.....	166	INLYTA ORAL TABLET 1 MG, 5 MG....	41
HEB INCONTROL ALCOHOL 70% PADS.....	166	INQOVI.....	84
HERCEPTIN HYLECTA.....	366	INREBIC.....	130
HERNEXEOS.....	403	INSULIN 1 ML SYRINGE.....	166
HUMIRA PEN.....	12	INSULIN SYR 0.3 ML 31GX1/4(1/2).....	166
HUMIRA PEN CROHNS-UC-HS START.	12	INSULIN SYR 0.5 ML 28G 12.7MM (OTC).....	166
HUMIRA PEN PSOR-UVEITS-ADOL HS	12	INSULIN SYRIN 0.5 ML 30GX1/2" (RX)	166
HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML.....	12	INSULIN SYRING 0.5 ML 27G 1/2"	166
HUMIRA(CF).....	12	INNER.....	166
HUMIRA(CF) PEDI CROHNS STARTER	12	INSULIN SYRINGE 0.3 ML.....	166
HUMIRA(CF) PEN.....	12	INSULIN SYRINGE 0.3 ML 31GX1/4.....	166
HUMIRA(CF) PEN CROHNS-UC-HS.....	12	INSULIN SYRINGE 0.5 ML.....	166
HUMIRA(CF) PEN PEDIATRIC UC.....	12	INSULIN SYRINGE 0.5 ML 31GX1/4.....	166
HUMIRA(CF) PEN PSOR-UV-ADOL HS.	12	INSULIN SYRINGE 1 ML.....	166
HYRNUO.....	309	INSULIN SYRINGE 1 ML 27G 1/2"	166
IBRANCE.....	253	INNER.....	166
IBTROZI.....	335	INSULIN SYRINGE 1 ML 27G 16MM....	166
<i>icatibant</i> .....	157	INSULIN SYRINGE 1 ML 28G 12.7MM (OTC).....	166
ICLUSIG.....	272	INSULIN SYRINGE 1 ML 30GX1/2"	166
IDHIFA.....	112	SHORT NEEDLE (OTC).....	166
<i>imatinib oral tablet 100 mg, 400 mg</i> .....	159	INSULIN SYRINGE 1 ML 31GX1/4".....	166
IMBRUVICA ORAL CAPSULE 140 MG, 70 MG.....	156	INSULIN SYRINGE NEEDLELESS.....	166
IMBRUVICA ORAL SUSPENSION.....	156	INSULIN SYRINGE-NEEDLE U-100 SYRINGE 0.3 ML 29 GAUGE, 1 ML 29 GAUGE X 1/2", 1/2 ML 28 GAUGE.....	166
IMBRUVICA ORAL TABLET.....	156	INSULIN U-500 SYRINGE-NEEDLE.....	166
IMDELLTRA.....	337	INSUPEN 30G ULTRAFIN NEEDLE.....	166
IMJUDO.....	368	INSUPEN 31G ULTRAFIN NEEDLE.....	166
IMKELDI.....	160	INSUPEN 32G 8MM PEN NEEDLE.....	166
IMPAVIDO.....	223	INSUPEN PEN NEEDLE 29GX12MM....	166
INCONTROL PEN NEEDLE 12MM 29G	166	INSUPEN PEN NEEDLE 31G 8MM.....	166
INCONTROL PEN NEEDLE 4MM 32G..	166	INSUPEN PEN NEEDLE 31GX3/16".....	166
INCONTROL PEN NEEDLE 5MM 31G..	166	INSUPEN PEN NEEDLE 32G 6MM (RX) .....	166
INCONTROL PEN NEEDLE 6MM 31G..	166	INSUPEN PEN NEEDLE 32GX4MM.....	166
INCONTROL PEN NEEDLE 8MM 31G..	166	INSUPEN PEN NEEDLE 33GX4MM.....	166
INCRELEX.....	215	ITOVEBI ORAL TABLET 3 MG, 9 MG..	163
<i>infliximab</i> .....	164	IV ANTISEPTIC WIPES.....	166
		IWILFIN.....	102
		JAKAFI.....	301
		<i>javygtor oral tablet,soluble</i> .....	302

JAYPIRCA ORAL TABLET 100 MG, 50 MG.....	270	LITE TOUCH PEN NEEDLE 29G.....	166
JEMPERLI.....	95	LITE TOUCH PEN NEEDLE 31G.....	166
JYNARQUE ORAL TABLET.....	359	LITETOUCH INS 0.3 ML 29GX1/2".....	166
KALYDECO.....	182	LITETOUCH INS 0.3 ML 30GX5/16".....	166
KENDALL ALCOHOL 70% PREP PAD.....	166	LITETOUCH INS 0.3 ML 31GX5/16".....	166
KERENDIA.....	136	LITETOUCH INS 0.5 ML 31GX5/16".....	166
KESIMPTA PEN.....	243	LITETOUCH SYR 0.5 ML 28GX1/2".....	166
KEYTRUDA.....	261	LITETOUCH SYR 0.5 ML 29GX1/2".....	166
KEYTRUDA QLEX.....	262	LITETOUCH SYR 0.5 ML 30GX5/16".....	166
KIMMTRAK.....	339	LITETOUCH SYRIN 1 ML 28GX1/2".....	166
KINERET.....	24	LITETOUCH SYRIN 1 ML 29GX1/2".....	166
KISQALI 200 MG DAILY DOSE.....	284	LITETOUCH SYRIN 1 ML 30GX5/16".....	166
KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG.....	285	LIVTENCITY.....	213
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3).....	284	LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG.....	370
KOSELUGO ORAL CAPSULE 10 MG, 25 MG.....	308	LOQTORZI.....	361
KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG, 7.5 MG.....	308	LORBRENA ORAL TABLET 100 MG, 25 MG.....	208
KRAZATI.....	11	LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG.....	327
KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG.....	28	LUNSUMIO.....	227
<i>lanreotide subcutaneous syringe 120 mg/0.5 ml</i> .....	186	LUPRON DEPOT.....	198
<i>lapatinib</i> .....	187	LUPRON DEPOT (3 MONTH).....	198
LAZCLUZE ORAL TABLET 240 MG, 80 MG.....	189	LUPRON DEPOT (4 MONTH).....	198
<i>lenalidomide</i> .....	191	LUPRON DEPOT (6 MONTH).....	198
LENVIMA.....	192	LUPRON DEPOT-PED (3 MONTH).....	200
<i>leuprolide acetate (3 month)</i> .....	195	LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT.....	200
<i>leuprolide subcutaneous kit</i> .....	194	LUTRATE DEPOT (3 MONTH).....	195
<i>lidocaine topical adhesive patch, medicated 5 %</i> .....	204	LYNOZYFIC INTRAVENOUS SOLUTION 2 MG/ML, 20 MG/ML.....	206
<i>lidocaine topical ointment</i> .....	203	LYNPARZA.....	244
<i>lidocaine-prilocaine topical cream</i> .....	205	LYTGObi.....	141
<i>lidocan iii</i> .....	204	MAGELLAN INSUL SYRINGE 0.3 ML.....	166
LISCO SPONGES 100/BAG.....	166	MAGELLAN INSUL SYRINGE 0.5 ML.....	166
LITE TOUCH 31GX1/4" PEN NEEDLE.....	166	MAGELLAN INSULIN SYR 0.3 ML.....	166
LITE TOUCH INSULIN 0.5 ML SYR.....	166	MAGELLAN INSULIN SYR 0.5 ML.....	166
LITE TOUCH INSULIN 1 ML SYR.....	166	MAGELLAN INSULIN SYRINGE 1 ML.....	166
LITE TOUCH INSULIN SYR 1 ML.....	166	MARGENZA.....	212
		MAVENCLAD (10 TABLET PACK).....	70
		MAVENCLAD (4 TABLET PACK).....	70
		MAVENCLAD (5 TABLET PACK).....	70
		MAVENCLAD (6 TABLET PACK).....	70
		MAVENCLAD (7 TABLET PACK).....	70
		MAVENCLAD (8 TABLET PACK).....	70
		MAVENCLAD (9 TABLET PACK).....	70

MAXICOMFORT II PEN NDL 31GX6MM.....	166	MONOJECT INSUL SYR U100 1 ML 3'S, 29GX1/2" (OTC).....	166
MAXICOMFORT INS 0.5 ML 27GX1/2"	166	MONOJECT INSUL SYR U100 1 ML W/O NEEDLE (OTC).....	166
MAXI-COMFORT INS 0.5 ML 28G.....	166	MONOJECT INSULIN SYR 0.3 ML.....	166
MAXICOMFORT INS 1 ML 27GX1/2" ..	166	MONOJECT INSULIN SYR 0.3 ML (OTC).....	166
MAXI-COMFORT INS 1 ML 28GX1/2" ..	166	MONOJECT INSULIN SYR 0.5 ML.....	166
MAXICOMFORT PEN NDL 29G X 5MM .....	166	MONOJECT INSULIN SYR 0.5 ML (OTC).....	166
MAXICOMFORT PEN NDL 29G X 8MM .....	166	MONOJECT INSULIN SYR 1 ML 3'S (OTC).....	166
MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG.....	312	MONOJECT INSULIN SYR U-100.....	166
MAYZENT STARTER(FOR 1MG MAINT).....	312	MONOJECT SYRINGE 0.3 ML.....	166
MAYZENT STARTER(FOR 2MG MAINT).....	312	MONOJECT SYRINGE 0.5 ML.....	166
MEKINIST ORAL RECON SOLN.....	363	MONOJECT SYRINGE 1 ML.....	166
MEKINIST ORAL TABLET 0.5 MG, 2 MG.....	364	<i>morphine concentrate oral solution</i> .....	155
MEKTOVI.....	54	MOUNJARO.....	151
<i>metirosine</i> .....	220	MS INSULIN SYR 1 ML 31GX5/16" (OTC).....	166
MICRODOT PEN NEEDLE 31GX6MM..	166	MS INSULIN SYRINGE 0.3 ML.....	166
MICRODOT PEN NEEDLE 32GX4MM..	166	NANO 2 GEN PEN NEEDLE 32G 4MM..	166
MICRODOT PEN NEEDLE 33GX4MM..	166	NANO PEN NEEDLE 32G 4MM.....	166
MICRODOT READYGARD NDL 31G 5MM OUTER.....	166	NATPARA.....	254
<i>mifepristone oral tablet 300 mg</i> .....	222	NERLYNX.....	230
MINI PEN NEEDLE 32G 5MM.....	166	NIKTIMVO.....	40
MINI PEN NEEDLE 32G 8MM.....	166	NINLARO.....	184
MINI PEN NEEDLE 33G 4MM.....	166	<i>nitisinone</i> .....	238
MINI PEN NEEDLE 33G 5MM.....	166	NIVESTYM.....	135
MINI PEN NEEDLE 33G 6MM.....	166	NORDITROPIN FLEXPPO.....	320
MINI ULTRA-THIN II PEN NDL 31G STERILE.....	166	NOVOFINE 30.....	166
MIPLYFFA.....	31	NOVOFINE 32G NEEDLES.....	166
<i>modafinil oral tablet 100 mg, 200 mg</i> .....	228	NOVOFINE PLUS PEN NDL 32GX1/6" ..	166
MODEYSO.....	94	NOVOTWIST.....	166
MONOJECT 0.5 ML SYRN 28GX1/2" .....	166	NUBEQA.....	81
MONOJECT 1 ML SYRN 27X1/2" .....	166	NUCALA SUBCUTANEOUS AUTO- INJECTOR.....	217
MONOJECT 1 ML SYRN 28GX1/2" (OTC).....	166	NUCALA SUBCUTANEOUS RECON SOLN.....	217
MONOJECT INSUL SYR U100 (OTC)....	166	NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML.....	217
MONOJECT INSUL SYR U100 .5ML,29GX1/2" (OTC).....	166	NUPLAZID.....	267
MONOJECT INSUL SYR U100 0.5 ML CONVERTS TO 29G (OTC).....	166	NURTEC ODT.....	289
MONOJECT INSUL SYR U100 1 ML.....	166	NYVEPRIA.....	257
		ODOMZO.....	323
		OFEV.....	233
		OGIVRI.....	365

OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG.....	237	PEN NEEDLES 8MM 31G 31GX8MM,STRL,SHORT (OTC).....	166
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION.....	362	<i>penicillamine oral tablet</i> .....	264
OJEMDA ORAL TABLET.....	362	PENTIPS PEN NEEDLE 29G 1/2".....	166
OJJAARA.....	226	PENTIPS PEN NEEDLE 31G 1/4".....	166
ONAPGO.....	27	PENTIPS PEN NEEDLE 31GX3/16" MINI, 5MM.....	166
ONUREG.....	42	PENTIPS PEN NEEDLE 31GX5/16" SHORT, 8MM.....	166
OPDIVO.....	239	PENTIPS PEN NEEDLE 32G 1/4".....	166
OPDIVO QVANTIG.....	240	PENTIPS PEN NEEDLE 32GX5/32" 4MM.....	166
OPDUALAG.....	241	PIP PEN NEEDLE 31G X 5MM.....	166
OPSUMIT.....	211	PIP PEN NEEDLE 32G X 4MM.....	166
ORENCIA.....	4	PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2).....	20
ORENCIA (WITH MALTOSE).....	2	<i>pirfenidone oral capsule</i> .....	268
ORENCIA CLICKJECT.....	4	<i>pirfenidone oral tablet 267 mg, 534 mg, 801 mg</i> .....	268
ORFADIN ORAL SUSPENSION.....	238	PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML.....	178
ORGOVYX.....	279	PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML.....	178
ORLISSA ORAL TABLET 150 MG, 200 MG.....	104	POMALYST.....	271
ORKAMBI ORAL TABLET.....	210	<i>posaconazole oral tablet, delayed release (dr/ec)</i> .....	273
ORSERDU ORAL TABLET 345 MG, 86 MG.....	103	PREFPLS INS SYR 1 ML 30GX5/16" (OTC).....	166
OSENVELT.....	87	PREVENT PEN NEEDLE 31GX1/4".....	166
OTEZLA.....	29	PREVENT PEN NEEDLE 31GX5/16".....	166
OTEZLA STARTER.....	29	PREVYMIS ORAL TABLET.....	193
OTEZLA XR.....	29	PRO COMFORT 0.5 ML 30GX1/2".....	166
<i>oxandrolone</i> .....	251	PRO COMFORT 0.5 ML 30GX5/16".....	166
OZEMPIC.....	150	PRO COMFORT 0.5 ML 31GX5/16".....	166
<i>pazopanib oral tablet 200 mg, 400 mg</i> .....	256	PRO COMFORT 1 ML 30GX1/2".....	166
PC UNIFINE PENTIPS 8MM NEEDLE SHORT.....	166	PRO COMFORT 1 ML 30GX5/16".....	166
PEGASYS.....	259	PRO COMFORT 1 ML 31GX5/16".....	166
PEMAZYRE.....	263	PRO COMFORT ALCOHOL 70% PADS.....	166
PEN NEEDLE 30G 5MM OUTER.....	166	PRO COMFORT PEN NDL 32G 8MM....	166
PEN NEEDLE 30G 8MM INNER.....	166	PRO COMFORT PEN NDL 32G X 1/4" ...	166
PEN NEEDLE 30G X 5/16".....	166	PRO COMFORT PEN NDL 4MM 32G....	166
PEN NEEDLE 31G X 1/4" HRI.....	166	PRO COMFORT PEN NDL 5MM 32G....	166
PEN NEEDLE 6MM 31G 6MM.....	166	PRODIGY INS SYR 1 ML 28GX1/2".....	166
PEN NEEDLE, DIABETIC NEEDLE 29 GAUGE X 1/2".....	166		
PEN NEEDLES 12MM 29G 29GX12MM,STRL.....	166		
PEN NEEDLES 4MM 32G.....	166		
PEN NEEDLES 5MM 31G 31GX5MM,STRL,MINI (OTC).....	166		

PRODIGY SYRNG 0.5 ML 31GX5/16"...	166	RUBRACA.....	300
PRODIGY SYRNGE 0.3 ML 31GX5/16" .	166	RYBELSUS.....	150
PURE CMFT SFTY PEN NDL 31G 5MM	166	RYBREVANT.....	23
PURE CMFT SFTY PEN NDL 31G 6MM	166	RYBREVANT FASPRO.....	22
PURE CMFT SFTY PEN NDL 32G 4MM	166	RYDAPT.....	221
PURE COMFORT ALCOHOL 70%		RYTELO.....	161
PADS.....	166	SAFESNAP INS SYR UNITS-100 0.3 ML	
PURE COMFORT PEN NDL 32G 4MM..	166	30GX5/16",10X10.....	166
PURE COMFORT PEN NDL 32G 5MM..	166	SAFESNAP INS SYR UNITS-100 0.5 ML	
PURE COMFORT PEN NDL 32G 6MM..	166	29GX1/2",10X10.....	166
PURE COMFORT PEN NDL 32G 8MM..	166	SAFESNAP INS SYR UNITS-100 0.5 ML	
<i>pyrimethamine</i> .....	275	30GX5/16",10X10.....	166
QINLOCK.....	293	SAFESNAP INS SYR UNITS-100 1 ML	
<i>quinine sulfate</i> .....	276	28GX1/2",10X10.....	166
QULIPTA.....	36	SAFESNAP INS SYR UNITS-100 1 ML	
RALDESY.....	367	29GX1/2",10X10.....	166
RAYA SURE PEN NEEDLE 29G 12MM.	166	SAFETY PEN NEEDLE 31G 4MM.....	166
RAYA SURE PEN NEEDLE 31G 4MM..	166	SAFETY PEN NEEDLE 5MM X 31G.....	166
RAYA SURE PEN NEEDLE 31G 5MM..	166	SAFETY SYRINGE 0.5 ML 30G 1/2".....	166
RAYA SURE PEN NEEDLE 31G 6MM..	166	<i>sapropterin oral tablet,soluble</i> .....	302
RELION INS SYR 0.3 ML 31GX6MM....	166	SCEMBLIX ORAL TABLET 100 MG, 20	
RELION INS SYR 0.5 ML 31GX6MM....	166	MG, 40 MG.....	32
RELION INS SYR 1 ML 31GX15/64".....	166	SECURESAFE PEN NDL 30GX5/16"	
RELI-ON INSULIN 1 ML SYR.....	166	OUTER.....	166
RETACRIT INJECTION SOLUTION		SECURESAFE SYR 0.5 ML 29G 1/2"	
10,000 UNIT/ML, 2,000 UNIT/ML,		OUTER.....	166
20,000 UNIT/2 ML, 20,000 UNIT/ML,		SECURESAFE SYRNG 1 ML 29G 1/2"	
3,000 UNIT/ML, 4,000 UNIT/ML, 40,000		OUTER.....	166
UNIT/ML.....	119	SELARSDI.....	378, 380
RETEVMO ORAL CAPSULE 40 MG, 80		SEROSTIM SUBCUTANEOUS RECON	
MG.....	307	SOLN 4 MG, 5 MG, 6 MG.....	322
RETEVMO ORAL TABLET 120 MG, 160		SIGNIFOR.....	255
MG, 40 MG, 80 MG.....	307	<i>sildenafil (pulm.hypertension) oral tablet..</i>	310
REVCIVI.....	106	SIRTURO.....	44
REVUFORJ ORAL TABLET 110 MG,		SKY SAFETY PEN NEEDLE 30G 5MM.	166
160 MG, 25 MG.....	283	SKY SAFETY PEN NEEDLE 30G 8MM.	166
REZDIFFRA.....	281	SKYRIZI.....	294
REZLIDHIA.....	245	SM ULT CFT 0.3 ML 31GX5/16(1/2).....	166
REZUROCK.....	46	<i>sodium oxybate</i> .....	314
RINVOQ.....	374	SOMATULINE DEPOT	
RINVOQ LQ.....	374	SUBCUTANEOUS SYRINGE 60 MG/0.2	
RITUXAN HYCELA.....	296	ML, 90 MG/0.3 ML.....	186
ROMVIMZA.....	393	SOMAVERT.....	260
ROZLYTREK ORAL CAPSULE 100 MG,		<i>sorafenib</i> .....	324
200 MG.....	115	SPRAVATO.....	124
ROZLYTREK ORAL PELLETS IN		STERILE PADS 2" X 2".....	166
PACKET.....	116	STIVARGA.....	278

STRENSIQ.....	33	TECHLITE 0.3 ML 31GX6MM (1/2).....	166
SUBVENITE ORAL SUSPENSION.....	185	TECHLITE 0.3 ML 31GX8MM (1/2).....	166
<i>sunitinib malate</i> .....	329	TECHLITE 0.5 ML 30GX12MM (1/2).....	166
SURE CMFT SFTY PEN NDL 31G 6MM	166	TECHLITE 0.5 ML 30GX8MM (1/2).....	166
SURE CMFT SFTY PEN NDL 32G 4MM	166	TECHLITE 0.5 ML 31GX6MM (1/2).....	166
SURE COMFORT 0.5 ML SYRINGE.....	166	TECHLITE 0.5 ML 31GX8MM (1/2).....	166
SURE COMFORT 1 ML SYRINGE.....	166	TECHLITE INS SYR 1 ML 29GX12MM.	166
SURE COMFORT 3/10 ML SYRINGE....	166	TECHLITE INS SYR 1 ML 30GX12MM.	166
SURE COMFORT 3/10 ML SYRINGE		TECHLITE INS SYR 1 ML 31GX6MM...	166
INSULIN SYRINGE.....	166	TECHLITE INS SYR 1 ML 31GX8MM...	166
SURE COMFORT 30G PEN NEEDLE.....	166	TECHLITE PEN NEEDLE 29GX1/2".....	166
SURE COMFORT ALCOHOL PREP		TECHLITE PEN NEEDLE 29GX3/8".....	166
PADS.....	166	TECHLITE PEN NEEDLE 31GX1/4".....	166
SURE COMFORT INS 0.3 ML 31GX1/4.	166	TECHLITE PEN NEEDLE 31GX3/16"....	166
SURE COMFORT INS 0.5 ML 31GX1/4.	166	TECHLITE PEN NEEDLE 31GX5/16"....	166
SURE COMFORT INS 1 ML 31GX1/4"...	166	TECHLITE PEN NEEDLE 32GX1/4".....	166
SURE COMFORT PEN NDL 29GX1/2"		TECHLITE PEN NEEDLE 32GX5/16"....	166
12.7MM.....	166	TECHLITE PEN NEEDLE 32GX5/32"....	166
SURE COMFORT PEN NDL 31G 5MM..	166	TECHLITE PLUS PEN NDL 32G 4MM...	166
SURE COMFORT PEN NDL 31G 8MM..	166	TECVAYLI.....	340
SURE COMFORT PEN NDL 32G 4MM..	166	TEPMETKO.....	343
SURE COMFORT PEN NDL 32G 6MM..	166	<i>teriparatide subcutaneous pen injector 20</i>	
SURE-FINE PEN NEEDLES 12.7MM.....	166	<i>mcg/dose (560mcg/2.24ml)</i> .....	344
SURE-FINE PEN NEEDLES 5MM.....	166	TERUMO INS SYRINGE U100-1 ML.....	166
SURE-FINE PEN NEEDLES 8MM.....	166	TERUMO INS SYRINGE U100-1/2 ML..	166
SURE-JECT INSU SYR U100 0.3 ML.....	166	TERUMO INS SYRINGE U100-1/3 ML..	166
SURE-JECT INSU SYR U100 0.5 ML.....	166	TERUMO INS SYRNG U100-1/2 ML.....	166
SURE-JECT INSU SYR U100 1 ML.....	166	<i>testosterone cypionate</i> .....	346
SURE-JECT INSUL SYR U100 1 ML.....	166	<i>testosterone enanthate</i> .....	347
SURE-JECT INSULIN SYRINGE 1 ML..	166	<i>testosterone transdermal gel in metered-</i>	
SURE-PREP ALCOHOL PREP PADS.....	166	<i>dose pump 12.5 mg/ 1.25 gram (1 %),</i>	
SYMPAZAN.....	71	<i>20.25 mg/1.25 gram (1.62 %).....</i>	345
SYNRIBO.....	246	<i>testosterone transdermal gel in packet 1 %</i>	
TABRECTA.....	64	<i>(25 mg/2.5gram), 1 % (50 mg/5 gram).....</i>	345
<i>tadalafil oral tablet 2.5 mg, 5 mg</i> .....	331	<i>tetrabenazine</i> .....	348
TAFINLAR ORAL CAPSULE.....	77	TEVIMBRA.....	350
TAFINLAR ORAL TABLET FOR		THALOMID ORAL CAPSULE 100 MG,	
SUSPENSION.....	78	150 MG, 200 MG, 50 MG.....	349
TAGRISSE.....	250	THINPRO INS SYRIN U100-0.3 ML.....	166
TALVEY.....	336	THINPRO INS SYRIN U100-0.5 ML.....	166
TALZENNA.....	333	THINPRO INS SYRIN U100-1 ML.....	166
TASIGNA ORAL CAPSULE 150 MG,		TIBSOVO.....	183
200 MG, 50 MG.....	231	TIVDAK.....	351
TAVNEOS.....	37	<i>tolvaptan (polycys kidney dis) oral tablets,</i>	
TAZVERIK.....	338	<i>sequential</i> .....	359
TECHLITE 0.3 ML 29GX12MM (1/2).....	166	TOPCARE CLICKFINE 31G X 1/4".....	166
TECHLITE 0.3 ML 30GX8MM (1/2).....	166	TOPCARE CLICKFINE 31G X 5/16".....	166

TOPCARE ULTRA COMFORT SYRINGE.....	166	TRUE COMFORT PRO ALCOHOL PADS.....	166
<i>torpenz oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg.....</i>	127	TRUE COMFORT SFTY 1 ML 30G 1/2" ..	166
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION	371	TRUE COMFRT PRO 0.5 ML 30G 1/2" ...	166
TREMFYA INTRAVENOUS.....	153	TRUE COMFRT SFTY 1 ML 30G 5/16" ..	166
TREMFYA ONE-PRESS.....	153	TRUE COMFRT SFTY 1 ML 31G 5/16" ..	166
TREMFYA PEN INDUCTION PK(2PEN).....	153	TRUE COMFRT SFTY 1 ML 32G 5/16" ..	166
TREMFYA PEN SUBCUTANEOUS PEN INJECTOR 200 MG/2 ML.....	153	TRUEPLUS PEN NEEDLE 29GX1/2" .....	166
TREMFYA SUBCUTANEOUS SYRINGE.....	153	TRUEPLUS PEN NEEDLE 31G X 1/4" ...	166
<i>tretinoin topical cream.....</i>	360	TRUEPLUS PEN NEEDLE 31GX3/16" ...	166
<i>tridacaine ii.....</i>	204	TRUEPLUS PEN NEEDLE 31GX5/16" ..	166
<i>trientine oral capsule 250 mg.....</i>	369	TRUEPLUS PEN NEEDLE 32GX5/32" ...	166
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL.....	107	TRUEPLUS SYR 0.3 ML 29GX1/2" .....	166
TRIKAFTA ORAL TABLETS, SEQUENTIAL.....	107	TRUEPLUS SYR 0.3 ML 30GX5/16" .....	166
TRUE CMFRT PRO 0.5 ML 30G 5/16" ....	166	TRUEPLUS SYR 0.3 ML 31GX5/16" .....	166
TRUE CMFRT PRO 0.5 ML 31G 5/16" ....	166	TRUEPLUS SYR 0.5 ML 28GX1/2" .....	166
TRUE CMFRT PRO 0.5 ML 32G 5/16" ....	166	TRUEPLUS SYR 0.5 ML 29GX1/2" .....	166
TRUE CMFT SFTY PEN NDL 31G 5MM	166	TRUEPLUS SYR 0.5 ML 30GX5/16" .....	166
TRUE CMFT SFTY PEN NDL 31G 6MM	166	TRUEPLUS SYR 0.5 ML 31GX5/16" .....	166
TRUE CMFT SFTY PEN NDL 32G 4MM	166	TRUEPLUS SYR 1 ML 28GX1/2" .....	166
TRUE COMFORT 0.5 ML 30G 1/2" .....	166	TRUEPLUS SYR 1 ML 29GX1/2" .....	166
TRUE COMFORT 0.5 ML 30G 5/16" .....	166	TRUEPLUS SYR 1 ML 30GX5/16" .....	166
TRUE COMFORT 0.5 ML 31G 5/16" .....	166	TRUEPLUS SYR 1 ML 31GX5/16" .....	166
TRUE COMFORT 0.5 ML 31GX5/16" .....	166	TRULICITY .....	149
TRUE COMFORT 1 ML 31GX5/16" .....	166	TRUQAP.....	63
TRUE COMFORT ALCOHOL 70% PADS.....	166	TRUXIMA.....	297
TRUE COMFORT PEN NDL 31G 8MM..	166	TUKYSA ORAL TABLET 150 MG, 50 MG.....	372
TRUE COMFORT PEN NDL 31GX5MM	166	TURALIO .....	266
TRUE COMFORT PEN NDL 31GX6MM	166	TYENNE.....	353, 355
TRUE COMFORT PEN NDL 32G 5MM..	166	TYENNE AUTOINJECTOR.....	355
TRUE COMFORT PEN NDL 32G 6MM..	166	TYMLOS.....	1
TRUE COMFORT PEN NDL 32GX4MM	166	UBRELVY .....	373
TRUE COMFORT PEN NDL 33G 4MM..	166	UDENYCA ONBODY .....	258
TRUE COMFORT PEN NDL 33G 5MM..	166	ULTICAR INS 0.3 ML 31GX1/4(1/2).....	166
TRUE COMFORT PEN NDL 33G 6MM..	166	ULTICARE INS 1 ML 31GX1/4" .....	166
TRUE COMFORT PRO 1 ML 30G 1/2" ...	166	ULTICARE INS SYR 0.3 ML 30G 8MM.	166
TRUE COMFORT PRO 1 ML 30G 5/16".	166	ULTICARE INS SYR 0.3 ML 31G 6MM.	166
TRUE COMFORT PRO 1 ML 31G 5/16".	166	ULTICARE INS SYR 0.3 ML 31G 8MM.	166
TRUE COMFORT PRO 1 ML 32G 5/16".	166	ULTICARE INS SYR 0.5 ML 30G 8MM (OTC).....	166
		ULTICARE INS SYR 0.5 ML 31G 6MM.	166
		ULTICARE INS SYR 0.5 ML 31G 8MM (OTC).....	166
		ULTICARE INS SYR 1 ML 30GX1/2" .....	166
		ULTICARE PEN NEEDLE 31GX3/16" .....	166
		ULTICARE PEN NEEDLE 6MM 31G.....	166

ULTICARE PEN NEEDLE 8MM 31G.....	166	ULTRA FLO 0.3 ML 30G 1/2" (1/2).....	166
ULTICARE PEN NEEDLES 12MM 29G.	166	ULTRA FLO 0.3 ML 30G 5/16"(1/2).....	166
ULTICARE PEN NEEDLES 4MM 32G		ULTRA FLO 0.3 ML 31G 5/16"(1/2).....	166
MICRO, 32GX4MM.....	166	ULTRA FLO PEN NEEDLE 31G 5MM...	166
ULTICARE PEN NEEDLES 6MM 32G...	166	ULTRA FLO PEN NEEDLE 31G 8MM...	166
ULTICARE SAFE PEN NDL 30G 8MM..	166	ULTRA FLO PEN NEEDLE 32G 4MM...	166
ULTICARE SAFE PEN NDL 5MM 30G..	166	ULTRA FLO PEN NEEDLE 33G 4MM...	166
ULTICARE SAFETY 0.5 ML 29GX1/2		ULTRA FLO PEN NEEDLES 12MM 29G	
(RX).....	166	.....	166
ULTICARE SYR 0.3 ML 29G 12.7MM....	166	ULTRA FLO SYR 0.3 ML 29GX1/2" .....	166
ULTICARE SYR 0.3 ML 30GX1/2" .....	166	ULTRA FLO SYR 0.3 ML 30G 5/16" .....	166
ULTICARE SYR 0.3 ML 31GX5/16"		ULTRA FLO SYR 0.3 ML 31G 5/16" .....	166
SHORT NDL.....	166	ULTRA FLO SYR 0.5 ML 29G 1/2" .....	166
ULTICARE SYR 0.5 ML 30GX1/2" .....	166	ULTRA THIN PEN NDL 32G X 4MM.....	166
ULTICARE SYR 0.5 ML 31GX5/16"		ULTRACARE INS 0.3 ML 30GX5/16" ....	166
SHORT NDL.....	166	ULTRACARE INS 0.3 ML 31GX5/16" ....	166
ULTICARE SYR 1 ML 31GX5/16" .....	166	ULTRACARE INS 0.5 ML 30GX1/2" .....	166
ULTIGUARD SAFE 1 ML 30G 12.7MM.	166	ULTRACARE INS 0.5 ML 30GX5/16" ....	166
ULTIGUARD SAFE0.3 ML 30G 12.7MM		ULTRACARE INS 0.5 ML 31GX5/16" ....	166
.....	166	ULTRACARE INS 1 ML 30G X 5/16" .....	166
ULTIGUARD SAFE0.5 ML 30G 12.7MM		ULTRACARE INS 1 ML 30GX1/2" .....	166
.....	166	ULTRACARE INS 1 ML 31G X 5/16".....	166
ULTIGUARD SAFEPACK 1 ML 31G		ULTRACARE PEN NEEDLE 31GX1/4" ..	166
8MM.....	166	ULTRACARE PEN NEEDLE 31GX3/16" 166	
ULTIGUARD SAFEPACK 29G 12.7MM	166	ULTRACARE PEN NEEDLE 31GX5/16" 166	
ULTIGUARD SAFEPACK 31G 5MM.....	166	ULTRACARE PEN NEEDLE 32GX1/4" ..	166
ULTIGUARD SAFEPACK 31G 6MM.....	166	ULTRACARE PEN NEEDLE 32GX3/16" 166	
ULTIGUARD SAFEPACK 31G 8MM.....	166	ULTRACARE PEN NEEDLE 32GX5/32" 166	
ULTIGUARD SAFEPACK 32G 4MM.....	166	ULTRACARE PEN NEEDLE 33GX5/32" 166	
ULTIGUARD SAFEPACK 32G 6MM.....	166	ULTRA-FINE 0.3 ML 30G 12.7MM.....	166
ULTIGUARD SAFEPK 0.3 ML 31G		ULTRA-FINE 0.3 ML 31G 6MM (1/2).....	166
8MM.....	166	ULTRA-FINE 0.3 ML 31G 8MM (1/2).....	166
ULTIGUARD SAFEPK 0.5 ML 31G		ULTRA-FINE 0.5 ML 30G 12.7MM.....	166
8MM.....	166	ULTRA-FINE INS SYR 1 ML 31G 6MM	166
ULTILET ALCOHOL STERL SWAB.....	166	ULTRA-FINE INS SYR 1 ML 31G 8MM	166
ULTILET INSULIN SYRINGE 0.3 ML...	166	ULTRA-FINE PEN NDL 29G 12.7MM....	166
ULTILET INSULIN SYRINGE 0.5 ML...	166	ULTRA-FINE PEN NEEDLE 31G 5MM..	166
ULTILET INSULIN SYRINGE 1 ML.....	166	ULTRA-FINE PEN NEEDLE 31G 8MM..	166
ULTILET PEN NEEDLE.....	166	ULTRA-FINE PEN NEEDLE 32G 6MM..	166
ULTILET PEN NEEDLE 4MM 32G.....	166	ULTRA-FINE SYR 0.5 ML 31G 6MM.....	166
ULTRA COMFORT 0.3 ML SYRINGE...	166	ULTRA-FINE SYR 0.5 ML 31G 8MM.....	166
ULTRA COMFORT 0.5 ML 28GX1/2"		ULTRA-FINE SYR 1 ML 30G 12.7MM...	166
CONVERTS TO 29G.....	166	ULTRA-THIN II 1 ML 31GX5/16" .....	166
ULTRA COMFORT 0.5 ML 29GX1/2" ....	166	ULTRA-THIN II INS 0.3 ML 30G.....	166
ULTRA COMFORT 0.5 ML SYRINGE...	166	ULTRA-THIN II INS 0.3 ML 31G.....	166
ULTRA COMFORT 1 ML 31GX5/16" .....	166	ULTRA-THIN II INS 0.5 ML 29G.....	166
ULTRA COMFORT 1 ML SYRINGE.....	166	ULTRA-THIN II INS 0.5 ML 30G.....	166

ULTRA-THIN II INS 0.5 ML 31G.....	166	<i>ustekinumab-aauz</i> .....	376
ULTRA-THIN II INS SYR 1 ML 29G.....	166	VALCHLOR.....	216
ULTRA-THIN II INS SYR 1 ML 30G.....	166	VANFLYTA.....	277
ULTRA-THIN II PEN NDL 29GX1/2".....	166	VANISHPOINT 0.5 ML 30GX1/2" SY	
ULTRA-THIN II PEN NDL 31GX5/16".....	166	OUTER.....	166
UNIFINE OTC PEN NEEDLE 31G 5MM	166	VANISHPOINT INS 1 ML 30GX3/16"....	166
UNIFINE OTC PEN NEEDLE 32G 4MM	166	VANISHPOINT U-100 29X1/2 SYR.....	166
UNIFINE PEN NEEDLE 32G 4MM.....	166	VENCLEXTA ORAL TABLET 10 MG,	
UNIFINE PENTIPS 12MM 29G		100 MG, 50 MG.....	390
29GX12MM, STRL.....	166	VENCLEXTA STARTING PACK.....	390
UNIFINE PENTIPS 31GX3/16"		VEOZAH.....	133
31GX5MM,STRL,MINI.....	166	VERIFINE INS SYR 1 ML 29G 1/2".....	166
UNIFINE PENTIPS 32G 4MM.....	166	VERIFINE PEN NEEDLE 29G 12MM.....	166
UNIFINE PENTIPS 32GX1/4".....	166	VERIFINE PEN NEEDLE 31G 5MM.....	166
UNIFINE PENTIPS 33GX5/32".....	166	VERIFINE PEN NEEDLE 31G X 6MM...	166
UNIFINE PENTIPS 6MM 31G.....	166	VERIFINE PEN NEEDLE 31G X 8MM...	166
UNIFINE PENTIPS MAX 30GX3/16".....	166	VERIFINE PEN NEEDLE 32G 6MM.....	166
UNIFINE PENTIPS NEEDLES 29G.....	166	VERIFINE PEN NEEDLE 32G X 4MM...	166
UNIFINE PENTIPS PLUS 29GX1/2"		VERIFINE PEN NEEDLE 32G X 5MM...	166
12MM.....	166	VERIFINE PLUS PEN NDL 31G 5MM...	166
UNIFINE PENTIPS PLUS 30GX3/16"....	166	VERIFINE PLUS PEN NDL 31G 8MM...	166
UNIFINE PENTIPS PLUS 31GX1/4"		VERIFINE PLUS PEN NDL 32G 4MM...	166
ULTRA SHORT, 6MM.....	166	VERIFINE PLUS PEN NDL 32G 4MM-	
UNIFINE PENTIPS PLUS 31GX3/16"		SHARPS CONTAINER.....	166
MINI.....	166	VERIFINE SYRING 0.5 ML 29G 1/2".....	166
UNIFINE PENTIPS PLUS 31GX5/16"		VERIFINE SYRING 1 ML 31G 5/16".....	166
SHORT.....	166	VERIFINE SYRNG 0.3 ML 31G 5/16".....	166
UNIFINE PENTIPS PLUS 32GX5/32".....	166	VERIFINE SYRNG 0.5 ML 31G 5/16".....	166
UNIFINE PENTIPS PLUS 33GX5/32".....	166	VERQUVO.....	391
UNIFINE PROTECT 30G 5MM.....	166	VERSALON ALL PURPOSE SPONGE	
UNIFINE PROTECT 30G 8MM.....	166	25'S,N-STERILE,3PLY.....	166
UNIFINE PROTECT 32G 4MM.....	166	VERZENIO.....	6
UNIFINE SAFECONTROL 30G 5MM.....	166	<i>vigabatrin</i> .....	392
UNIFINE SAFECONTROL 30G 8MM.....	166	<i>vigadrone</i> .....	392
UNIFINE SAFECONTROL 31G 5MM.....	166	<i>vigpoder</i> .....	392
UNIFINE SAFECONTROL 31G 6MM.....	166	VITRAKVI ORAL CAPSULE 100 MG,	
UNIFINE SAFECONTROL 31G 8MM.....	166	25 MG.....	188
UNIFINE SAFECONTROL 32G 4MM.....	166	VITRAKVI ORAL SOLUTION.....	188
UNIFINE ULTRA PEN NDL 31G 5MM..	166	VIVIMUSTA.....	48
UNIFINE ULTRA PEN NDL 31G 6MM..	166	VIZIMPRO.....	79
UNIFINE ULTRA PEN NDL 31G 8MM..	166	VONJO.....	252
UNIFINE ULTRA PEN NDL 32G 4MM..	166	VOQUEZNA.....	395
UPTRAVI ORAL TABLET 1,000 MCG,		VORANIGO.....	397
1,200 MCG, 1,400 MCG, 1,600 MCG, 200		<i>voriconazole oral suspension for</i>	
MCG, 400 MCG, 600 MCG, 800 MCG.....	305	<i>reconstitution</i> .....	398
UPTRAVI ORAL TABLETS,DOSE		VOSEVI.....	318
PACK.....	305	VOWST.....	129

VUMERITY .....	93	ZURZUVAE ORAL CAPSULE 20 MG,	
VYALEV .....	139	25 MG, 30 MG.....	404
VYLOY .....	402	ZYDELIG .....	158
VYNDAMAX.....	332	ZYKADIA .....	66
WEBCOL ALCOHOL PREPS		ZYNLONTA.....	207
20'S,LARGE .....	166	ZYNYZ.....	282
WELIREG.....	47		
WINREVAIR.....	325		
XALKORI ORAL CAPSULE.....	75		
XALKORI ORAL PELLETT 150 MG, 20			
MG, 50 MG.....	76		
XDEMVI .....	209		
XELJANZ.....	357		
XELJANZ XR.....	357		
XERMELO.....	342		
XGEVA.....	88		
XIFAXAN ORAL TABLET 200 MG, 550			
MG.....	286		
XOLAIR.....	247		
XOSPATA.....	146		
XPOVIO ORAL TABLET 100			
MG/WEEK (50 MG X 2), 40 MG/WEEK			
(10 MG X 4), 40 MG/WEEK (40 MG X			
1), 40MG TWICE WEEK (40 MG X 2), 60			
MG/WEEK (60 MG X 1), 60MG TWICE			
WEEK (120 MG/WEEK), 80 MG/WEEK			
(40 MG X 2), 80 MG/WEEK (80 MG X			
1), 80MG TWICE WEEK (160			
MG/WEEK).....	306		
XTANDI ORAL CAPSULE.....	117		
XTANDI ORAL TABLET 40 MG, 80 MG			
.....	117		
YERVOY .....	180		
YESINTEK.....	382, 384		
YONSA.....	8		
YUFLYMA(CF).....	14		
YUFLYMA(CF) AI CROHN'S-UC-HS.....	14		
YUFLYMA(CF) AUTOINJECTOR.....	14		
ZEJULA ORAL CAPSULE.....	235		
ZEJULA ORAL TABLET .....	235		
ZELBORAF .....	389		
ZIIHERA.....	399		
ZIRABEV .....	52		
ZOLADEX.....	152		
ZTALMY .....	143		
ZTLIDO .....	204		